

Lifts and generalized vertices for Brauer characters of solvable groups

Mark L. Lewis

Kent State University

April 14, 2010

Ischia Group Theory Conference 2010

(Joint work with J. P. Cossey – University of Akron)

Lifts of Brauer Characters

- Throughout G will be a finite group and p will be a prime.

Lifts of Brauer Characters

- Throughout G will be a finite group and p will be a prime.
- $\text{IBr}(G)$ is the set of irreducible p -Brauer characters of G .

Lifts of Brauer Characters

- Throughout G will be a finite group and p will be a prime.
- $\text{IBr}(G)$ is the set of irreducible p -Brauer characters of G .
- G° denotes the p -regular elements of G . If χ is a character of G , then we use χ° to denote the restriction of χ to G° .

Lifts of Brauer Characters

- Throughout G will be a finite group and p will be a prime.
- $\text{IBr}(G)$ is the set of irreducible p -Brauer characters of G .
- G° denotes the p -regular elements of G . If χ is a character of G , then we use χ° to denote the restriction of χ to G° .

Definition

Given a p -Brauer character $\varphi \in \text{IBr}(G)$, we say $\chi \in \text{Irr}(G)$ is a *lift* of φ if $\chi^\circ = \varphi$.

Lifts of Brauer Characters

- Throughout G will be a finite group and p will be a prime.
- $\text{IBr}(G)$ is the set of irreducible p -Brauer characters of G .
- G° denotes the p -regular elements of G . If χ is a character of G , then we use χ° to denote the restriction of χ to G° .

Definition

Given a p -Brauer character $\varphi \in \text{IBr}(G)$, we say $\chi \in \text{Irr}(G)$ is a *lift* of φ if $\chi^\circ = \varphi$.

- When G is p -solvable, the Fong-Swan theorem implies that φ has a lift.

Lifts of Brauer Characters

- Much of the study of lifts has focused on particular canonical sets of lifts.

Lifts of Brauer Characters

- Much of the study of lifts has focused on particular canonical sets of lifts.
- J. P. Cossey has initiated the study of all lifts of φ . For example, when $|G|$ is odd, he has shown that the number of lifts of φ can be bounded in terms of a vertex for φ .

Lifts of Brauer Characters

- Much of the study of lifts has focused on particular canonical sets of lifts.
- J. P. Cossey has initiated the study of all lifts of φ . For example, when $|G|$ is odd, he has shown that the number of lifts of φ can be bounded in terms of a vertex for φ .
- We will show that the oddness hypothesis in Cossey's results can be removed in certain cases.

Vertices

- In a p -solvable group G , we say Q is a *vertex* for $\varphi \in \text{IBr}(G)$ if there is a subgroup U so that φ is induced from a p -Brauer character of U having p' -degree and Q is a Sylow p -subgroup of U .

Vertices

- In a p -solvable group G , we say Q is a *vertex* for $\varphi \in \text{IBr}(G)$ if there is a subgroup U so that φ is induced from a p -Brauer character of U having p' -degree and Q is a Sylow p -subgroup of U .
- It is known that all of the vertices for φ are conjugate in G .

Vertices

- In a p -solvable group G , we say Q is a *vertex* for $\varphi \in \text{IBr}(G)$ if there is a subgroup U so that φ is induced from a p -Brauer character of U having p' -degree and Q is a Sylow p -subgroup of U .
- It is known that all of the vertices for φ are conjugate in G .
- Cossey showed that if $|G|$ is odd and Q is a vertex for φ , then the number of lifts of φ is at most $|Q : Q'|$.

Vertices

We now remove the hypothesis that $|G|$ is odd.

However, we do need to add some hypotheses:

- 1 G is p -solvable
- 2 p is an odd prime
- 3 Q is abelian

Vertices

We now remove the hypothesis that $|G|$ is odd.

However, we do need to add some hypotheses:

- 1 G is p -solvable
- 2 p is an odd prime
- 3 Q is abelian

Theorem

Let G be a p -solvable group and let p be an odd prime. If $\varphi \in \text{IBr}(G)$ has abelian vertex Q , then the number of lifts of φ is at most $|Q|$.

Generalized vertices

We use the generalized vertices defined by Cossey. To do this, we need p -factored characters.

Generalized vertices

We use the generalized vertices defined by Cossey. To do this, we need p -factored characters.

- A character $\chi \in \text{Irr}(G)$ is p -factored if $\chi = \alpha\beta$ where α is p -special and β is p' -special.

Generalized vertices

We use the generalized vertices defined by Cossey. To do this, we need p -factored characters.

- A character $\chi \in \text{Irr}(G)$ is p -factored if $\chi = \alpha\beta$ where α is p -special and β is p' -special.
- Let $\chi \in \text{Irr}(G)$. Then (Q, δ) is a *generalized vertex* for χ if there is a subgroup U with a p -factored character $\psi \in \text{Irr}(U)$ and Sylow p -subgroup Q of U so that $\psi^G = \chi$ and δ is the restriction to Q of the p -special factor of ψ .

Generalized vertices

Since any primitive irreducible character of a p -solvable group is p -factored and p -special characters restrict irreducibly to a Sylow p -subgroup, all characters have generalized vertices.

However, for a general irreducible character χ , it seems unlikely that one can say anything useful about the set of all generalized vertices for χ .

Generalized vertices

Since any primitive irreducible character of a p -solvable group is p -factored and p -special characters restrict irreducibly to a Sylow p -subgroup, all characters have generalized vertices.

However, for a general irreducible character χ , it seems unlikely that one can say anything useful about the set of all generalized vertices for χ .

Theorem

(Cossey) Suppose $|G|$ is odd and $\chi \in \text{Irr}(G)$. Let (Q, δ) be a generalized vertex for χ . If $\chi^0 \in \text{IBr}(G)$, then

Generalized vertices

Since any primitive irreducible character of a p -solvable group is p -factored and p -special characters restrict irreducibly to a Sylow p -subgroup, all characters have generalized vertices.

However, for a general irreducible character χ , it seems unlikely that one can say anything useful about the set of all generalized vertices for χ .

Theorem

(Cossey) Suppose $|G|$ is odd and $\chi \in \text{Irr}(G)$. Let (Q, δ) be a generalized vertex for χ . If $\chi^0 \in \text{IBr}(G)$, then

- 1 δ is linear

Generalized vertices

Since any primitive irreducible character of a p -solvable group is p -factored and p -special characters restrict irreducibly to a Sylow p -subgroup, all characters have generalized vertices.

However, for a general irreducible character χ , it seems unlikely that one can say anything useful about the set of all generalized vertices for χ .

Theorem

(Cossey) Suppose $|G|$ is odd and $\chi \in \text{Irr}(G)$. Let (Q, δ) be a generalized vertex for χ . If $\chi^0 \in \text{IBr}(G)$, then

- 1 δ is linear
- 2 all generalized vertices for χ are conjugate to (Q, δ) .

Generalized vertices

Again, we will remove the hypothesis that $|G|$ is odd.

We do need to assume that p is odd.

Generalized vertices

Again, we will remove the hypothesis that $|G|$ is odd.

We do need to assume that p is odd.

To get δ linear, we use a recent theorem of Navarro:

Theorem

(Navarro) Let G be a p -solvable group for odd prime p . Let $\chi \in \text{Irr}(G)$ be p -special. If $\chi(1) > 1$, then χ° is not in $\text{IBr}(G)$.

Generalized vertices

Again, we will remove the hypothesis that $|G|$ is odd.

We do need to assume that p is odd.

To get δ linear, we use a recent theorem of Navarro:

Theorem

(Navarro) Let G be a p -solvable group for odd prime p . Let $\chi \in \text{Irr}(G)$ be p -special. If $\chi(1) > 1$, then χ° is not in $\text{IBr}(G)$.

Note: this theorem is not true if $p = 2$.

Generalized vertices

As a corollary to Navarro's result, we obtain the following:

Corollary

Let G be a p -solvable group where p is an odd prime. If $\chi \in \text{Irr}(G)$ satisfies $\chi^\circ \in \text{IBr}(G)$ and has generalized vertex (Q, δ) , then δ is linear.

Generalized vertices

As a corollary to Navarro's result, we obtain the following:

Corollary

Let G be a p -solvable group where p is an odd prime. If $\chi \in \text{Irr}(G)$ satisfies $\chi^\circ \in \text{IBr}(G)$ and has generalized vertex (Q, δ) , then δ is linear.

Notice that Q is now a vertex for χ° .

Generalized vertices

As a corollary to Navarro's result, we obtain the following:

Corollary

Let G be a p -solvable group where p is an odd prime. If $\chi \in \text{Irr}(G)$ satisfies $\chi^\circ \in \text{IBr}(G)$ and has generalized vertex (Q, δ) , then δ is linear.

Notice that Q is now a vertex for χ° .

If $p = 2$, this corollary is not true. In $\text{GL}_2(3)$, there is a counterexample.

Generalized vertices

We now prove:

Theorem

Let G be a p -solvable group and p an odd prime. If $\chi \in \text{Irr}(G)$ with $\chi^\circ \in \text{IBr}(G)$, then all the generalized vertices for χ are conjugate.

Generalized vertices

We now prove:

Theorem

Let G be a p -solvable group and p an odd prime. If $\chi \in \text{Irr}(G)$ with $\chi^\circ \in \text{IBr}(G)$, then all the generalized vertices for χ are conjugate.

When $p = 2$, the theorem is not true.

Counting lifts

Now, we return to our original question of counting the number of lifts of a given Brauer character $\varphi \in \text{IBr}(G)$.

Counting lifts

Now, we return to our original question of counting the number of lifts of a given Brauer character $\varphi \in \text{IBr}(G)$.

The main work is to count the number of lifts of φ with a given generalized vertex.

Counting lifts

Now, we return to our original question of counting the number of lifts of a given Brauer character $\varphi \in \text{IBr}(G)$.

The main work is to count the number of lifts of φ with a given generalized vertex.

Theorem

Assume that G is a p -solvable group and p is an odd prime. Suppose that $\varphi \in \text{IBr}(G)$ has vertex subgroup Q that is abelian, and let $\delta \in \text{Irr}(Q)$. Then $|L_\varphi(Q, \delta)| \leq |N_G(Q) : N_G(Q, \delta)|$.

Counting lifts

We use the count on the number of lifts of φ with a given generalized vertex to get the bound on the total number of lifts of φ :

Counting lifts

We use the count on the number of lifts of φ with a given generalized vertex to get the bound on the total number of lifts of φ :

Take $\delta_1, \dots, \delta_k$ to be representatives of the $N_G(Q)$ orbits of the characters of Q .

Counting lifts

We use the count on the number of lifts of φ with a given generalized vertex to get the bound on the total number of lifts of φ :

Take $\delta_1, \dots, \delta_k$ to be representatives of the $N_G(Q)$ orbits of the characters of Q .

One can show that every generalized vertex for a lift of φ is G -conjugate to (Q, δ_i) for some i . Thus, $|L_\varphi| = \sum_{i=1}^k |L_\varphi(Q, \delta_i)|$.

Counting lifts

We use the count on the number of lifts of φ with a given generalized vertex to get the bound on the total number of lifts of φ :

Take $\delta_1, \dots, \delta_k$ to be representatives of the $N_G(Q)$ orbits of the characters of Q .

One can show that every generalized vertex for a lift of φ is G -conjugate to (Q, δ_i) for some i . Thus, $|L_\varphi| = \sum_{i=1}^k |L_\varphi(Q, \delta_i)|$.

Applying the count for the generalized vertices:

$$\sum_{i=1}^k |L_\varphi(Q, \delta_i)| \leq \sum_{i=1}^k |N_G(Q) : N_G(Q, \delta_i)|.$$

Counting lifts

We use the count on the number of lifts of φ with a given generalized vertex to get the bound on the total number of lifts of φ :

Take $\delta_1, \dots, \delta_k$ to be representatives of the $N_G(Q)$ orbits of the characters of Q .

One can show that every generalized vertex for a lift of φ is G -conjugate to (Q, δ_i) for some i . Thus, $|L_\varphi| = \sum_{i=1}^k |L_\varphi(Q, \delta_i)|$.

Applying the count for the generalized vertices:

$$\sum_{i=1}^k |L_\varphi(Q, \delta_i)| \leq \sum_{i=1}^k |N_G(Q) : N_G(Q, \delta_i)|.$$

Finally, counting the sizes of the orbits of $N_G(Q)$ on the linear characters of Q , we obtain $\sum_{i=1}^k |N_G(Q) : N_G(Q, \delta_i)| = |Q|$.

Counting lifts

We use the count on the number of lifts of φ with a given generalized vertex to get the bound on the total number of lifts of φ :

Take $\delta_1, \dots, \delta_k$ to be representatives of the $N_G(Q)$ orbits of the characters of Q .

One can show that every generalized vertex for a lift of φ is G -conjugate to (Q, δ_i) for some i . Thus, $|L_\varphi| = \sum_{i=1}^k |L_\varphi(Q, \delta_i)|$.

Applying the count for the generalized vertices:

$$\sum_{i=1}^k |L_\varphi(Q, \delta_i)| \leq \sum_{i=1}^k |N_G(Q) : N_G(Q, \delta_i)|.$$

Finally, counting the sizes of the orbits of $N_G(Q)$ on the linear characters of Q , we obtain $\sum_{i=1}^k |N_G(Q) : N_G(Q, \delta_i)| = |Q|$.

Combining: $|L_\varphi| \leq |Q|$. (As desired.)

Q abelian?

We would like to remove the hypothesis that Q is abelian.

Q abelian?

We would like to remove the hypothesis that Q is abelian.

The point where Q is abelian is needed is in counting the number of lifts of φ with vertex (Q, δ) .

Q abelian?

We would like to remove the hypothesis that Q is abelian.

The point where Q is abelian is needed is in counting the number of lifts of φ with vertex (Q, δ) .

We review the proof of the number of lifts with a given vertex. Our proof is an inductive argument.

Q abelian?

We would like to remove the hypothesis that Q is abelian.

The point where Q is abelian is needed is in counting the number of lifts of φ with vertex (Q, δ) .

We review the proof of the number of lifts with a given vertex. Our proof is an inductive argument.

Let $\chi \in \text{Irr}(G)$ be a lift of φ with vertex (Q, δ) . Let N be maximal so that N is normal in G and the irreducible constituents of χ_N are factored.

Q abelian?

We would like to remove the hypothesis that Q is abelian.

The point where Q is abelian is needed is in counting the number of lifts of φ with vertex (Q, δ) .

We review the proof of the number of lifts with a given vertex. Our proof is an inductive argument.

Let $\chi \in \text{Irr}(G)$ be a lift of φ with vertex (Q, δ) . Let N be maximal so that N is normal in G and the irreducible constituents of χ_N are factored.

Let $\alpha\beta$ be an irreducible constituent of χ_N where α is p' -special and β is p -special.

Q abelian?

We first show that we may assume that α is G -invariant using the inductive hypothesis.

Q abelian?

We first show that we may assume that α is G -invariant using the inductive hypothesis.

Let T be the stabilizer of α in G .

Q abelian?

We first show that we may assume that α is G -invariant using the inductive hypothesis.

Let T be the stabilizer of α in G .

There is a unique Brauer character η on T so that $\eta^G = \varphi$ and α° is a constituent of η_N .

Q abelian?

We first show that we may assume that α is G -invariant using the inductive hypothesis.

Let T be the stabilizer of α in G .

There is a unique Brauer character η on T so that $\eta^G = \varphi$ and α° is a constituent of η_N .

By replacing α with a G -conjugate if necessary, we may assume that $Q \leq T$ and Q is a vertex for η .

Q abelian?

We first show that we may assume that α is G -invariant using the inductive hypothesis.

Let T be the stabilizer of α in G .

There is a unique Brauer character η on T so that $\eta^G = \varphi$ and α° is a constituent of η_N .

By replacing α with a G -conjugate if necessary, we may assume that $Q \leq T$ and Q is a vertex for η .

Let $\delta_1, \dots, \delta_l$ be representatives for the orbits of the action $N_T(Q)$ on the $N_G(Q)$ -orbit of δ .

Q abelian?

It is not difficult to see that there is a bijection between $\bigcup L_{\eta}(Q, \delta_i)$ and $L_{\varphi}(Q, \delta)$.

Q abelian?

It is not difficult to see that there is a bijection between $\bigcup L_\eta(Q, \delta_i)$ and $L_\varphi(Q, \delta)$.

So, $|L_\varphi(Q, \delta)| = \sum_{i=1}^n |L_\eta(Q, \delta_i)|$.

Q abelian?

It is not difficult to see that there is a bijection between $\bigcup L_\eta(Q, \delta_i)$ and $L_\varphi(Q, \delta)$.

So, $|L_\varphi(Q, \delta)| = \sum_{i=1}^n |L_\eta(Q, \delta_i)|$.

If $T < G$, then $|L_\eta(Q, \delta_i)| \leq |N_T(Q) : N_T(Q, \delta_i)|$ for all i .

Q abelian?

It is not difficult to see that there is a bijection between $\bigcup L_\eta(Q, \delta_i)$ and $L_\varphi(Q, \delta)$.

$$\text{So, } |L_\varphi(Q, \delta)| = \sum_{i=1}^n |L_\eta(Q, \delta_i)|.$$

If $T < G$, then $|L_\eta(Q, \delta_i)| \leq |N_T(Q) : N_T(Q, \delta_i)|$ for all i .

$$\text{Also, } \sum_{i=1}^n |N_T(Q) : N_T(Q, \delta_i)| = |N_G(Q) : N_G(Q, \delta)|.$$

Q abelian?

It is not difficult to see that there is a bijection between $\bigcup L_\eta(Q, \delta_i)$ and $L_\varphi(Q, \delta)$.

So, $|L_\varphi(Q, \delta)| = \sum_{i=1}^n |L_\eta(Q, \delta_i)|$.

If $T < G$, then $|L_\eta(Q, \delta_i)| \leq |N_T(Q) : N_T(Q, \delta_i)|$ for all i .

Also, $\sum_{i=1}^n |N_T(Q) : N_T(Q, \delta_i)| = |N_G(Q) : N_G(Q, \delta)|$.

Combining, we obtain: $|L_\varphi(Q, \delta)| \leq |N_G(Q) : N_G(Q, \delta)|$.

Q abelian?

It is not difficult to see that there is a bijection between $\bigcup L_\eta(Q, \delta_i)$ and $L_\varphi(Q, \delta)$.

$$\text{So, } |L_\varphi(Q, \delta)| = \sum_{i=1}^n |L_\eta(Q, \delta_i)|.$$

If $T < G$, then $|L_\eta(Q, \delta_i)| \leq |N_T(Q) : N_T(Q, \delta_i)|$ for all i .

$$\text{Also, } \sum_{i=1}^n |N_T(Q) : N_T(Q, \delta_i)| = |N_G(Q) : N_G(Q, \delta)|.$$

Combining, we obtain: $|L_\varphi(Q, \delta)| \leq |N_G(Q) : N_G(Q, \delta)|$.

Thus, $T = G$ and α is G -invariant.

Q abelian?

However, when we go to use the inductive hypothesis on β , things do not work as well.

Q abelian?

However, when we go to use the inductive hypothesis on β , things do not work as well.

Suppose I , the stabilizer of β in G , is chosen so that $Q \leq I$ and ζ_1, \dots, ζ_m are the characters in $\text{IBr}(I)$ with vertex Q that induce φ .

Q abelian?

However, when we go to use the inductive hypothesis on β , things do not work as well.

Suppose I , the stabilizer of β in G , is chosen so that $Q \leq I$ and ζ_1, \dots, ζ_m are the characters in $\text{IBr}(I)$ with vertex Q that induce φ .

We can show that $|L_\varphi(Q, \delta)|$ equals $\sum_{i=1}^m |L_{\zeta_i}(Q, \delta)|$.

Q abelian?

If $I < G$, we can use the inductive hypothesis to see that the result holds for each ζ_j .

Q abelian?

If $I < G$, we can use the inductive hypothesis to see that the result holds for each ζ_j .

That is, $|L_{\zeta_j}(Q, \delta)|$ is at most $|\mathbf{N}_I(Q) : \mathbf{N}_I(Q, \delta)|$.

Q abelian?

If $I < G$, we can use the inductive hypothesis to see that the result holds for each ζ_j .

That is, $|L_{\zeta_j}(Q, \delta)|$ is at most $|N_I(Q) : N_I(Q, \delta)|$.

With this, we deduce $|L_\varphi(Q, \delta)|$ is at most $m|N_I(Q) : N_I(Q, \delta)|$.

Q abelian?

If $I < G$, we can use the inductive hypothesis to see that the result holds for each ζ_j .

That is, $|L_{\zeta_j}(Q, \delta)|$ is at most $|N_I(Q) : N_I(Q, \delta)|$.

With this, we deduce $|L_\varphi(Q, \delta)|$ is at most $m|N_I(Q) : N_I(Q, \delta)|$.

To prove the result, we need $m \leq |N_G(Q) : N_I(Q)|$.

Q abelian?

When Q is abelian, we can prove that $Q \leq N$.

Q abelian?

When Q is abelian, we can prove that $Q \leq N$.

We then have the Frattini argument available to see that $G = NN_G(Q)$.

Q abelian?

When Q is abelian, we can prove that $Q \leq N$.

We then have the Frattini argument available to see that $G = NN_G(Q)$.

It follows $|G : I| = |N_G(Q) : N_I(Q)|$.

Q abelian?

When Q is abelian, we can prove that $Q \leq N$.

We then have the Frattini argument available to see that $G = NN_G(Q)$.

It follows $|G : I| = |N_G(Q) : N_I(Q)|$.

Also, $\varphi(1) = |G : I|\zeta_1(1)$ and $\varphi(1) \geq \sum_{i=1}^m \zeta_i(1) = m\zeta_1(1)$.

Q abelian?

When Q is abelian, we can prove that $Q \leq N$.

We then have the Frattini argument available to see that $G = NN_G(Q)$.

It follows $|G : I| = |N_G(Q) : N_I(Q)|$.

Also, $\varphi(1) = |G : I|\zeta_1(1)$ and $\varphi(1) \geq \sum_{i=1}^m \zeta_i(1) = m\zeta_1(1)$.

We conclude that $m \leq |G : I|$.

Q abelian?

Thus, when Q is abelian, we may assume β is G -invariant.

Q abelian?

Thus, when Q is abelian, we may assume β is G -invariant.

Now, α and β both G -invariant implies $N = G$.

Q abelian?

Thus, when Q is abelian, we may assume β is G -invariant.

Now, α and β both G -invariant implies $N = G$.

The result is immediate when $G = N$.

Q abelian?

Thus, when Q is abelian, we may assume β is G -invariant.

Now, α and β both G -invariant implies $N = G$.

The result is immediate when $G = N$.

When Q is not abelian, there is no reason to believe that $Q \leq N$.

An open question

Hence, we ask the follow question:

An open question

Hence, we ask the follow question:

Question:

Let G be a p -solvable group. Suppose $\varphi \in \text{IBr}(G)$ has vertex Q . Suppose $Q \leq I \leq G$. Is it true that the number of characters in $\text{IBr}(I)$ with vertex Q that induce φ is at most $|\text{N}_G(Q) : \text{N}_I(Q)|$?

An open question

Hence, we ask the follow question:

Question:

Let G be a p -solvable group. Suppose $\varphi \in \text{IBr}(G)$ has vertex Q . Suppose $Q \leq I \leq G$. Is it true that the number of characters in $\text{IBr}(I)$ with vertex Q that induce φ is at most $|\text{N}_G(Q) : \text{N}_I(Q)|$?

If the answer is yes, when p is odd, then we can remove the hypothesis that Q is abelian.

An open question

Hence, we ask the follow question:

Question:

Let G be a p -solvable group. Suppose $\varphi \in \text{IBr}(G)$ has vertex Q . Suppose $Q \leq I \leq G$. Is it true that the number of characters in $\text{IBr}(I)$ with vertex Q that induce φ is at most $|\text{N}_G(Q) : \text{N}_I(Q)|$?

If the answer is yes, when p is odd, then we can remove the hypothesis that Q is abelian.

We have not been able to settle this question at this time.

An open question

Interestingly, the question does have a positive answer when $|G|$ is odd or when $p = 2$.

An open question

Interestingly, the question does have a positive answer when $|G|$ is odd or when $p = 2$.

Also, when p is odd, we can prove that if G is a minimal counterexample, then I is a maximal subgroup, $|G : I|$ is a power of 2, and φ restricts homogeneously to every normal subgroup of G contained in I . Furthermore, writing N for the core of I in G and M for a normal subgroup of G so that M/N is a chief factor of G , if α is the irreducible constituent of φ_N , then α^M has a unique irreducible constituent.