

Dedicated to the memory of

Michio Suzuki

Dedicated to the memory of **Michio Suzuki**




Chiba (Japan), 2 October 1926 - Mitaka (Japan), 31 May **1998**



Michio Suzuki

www-history.mcs.st-andrews.ac.uk/Biographies/Suzuki-Michio.html


<https://en.wikipedia.org/wiki/Michio-Suzuki>




Structure of a Group and the Structure of its Lattice of Subgroups


Ergebnisse der Mathematik und ihrer Grenzgebiete

Springer Verlag, 1956



Group Theory I
*Grundlehren der mathematischen
Wissenschaften*
Springer Verlag, 1982



 **Group Theory II**
*Grundlehren der mathematischen
Wissenschaften*
Springer Verlag, 1986

Michio Suzuki (1926–1998)

Michael Aschbacher, Helmut Bender, Walter Feit,
and Ronald Solomon

Editor's Note. Michio Suzuki, an early leader in the effort to classify finite simple groups, died Mar. 31, 1998, in Tokyo at the age of seventy-one. Born October 2, 1926, in Japan, he obtained his Ph.D. from the University of Tokyo in 1952, with Shunji Uryu as his advisor. Suzuki's teachers included Shiro Aronson and Kazuhiko Hosokawa. Suzuki assumed a faculty position at the University of Illinois, Urbana-Champaign, beginning the next year. In 1956–57 he took a leave of absence to work at Harvard University as research associate with Richard Brauer, with support from the National Science Foundation. He was a professor in the Center for Advanced Study of the University of Illinois from 1960 until his death.

Suzuki held a postdoctoral fellowship in 1952–53 and a Guggenheim Fellowship in 1962–63, received the Academy Prize from the Japan Academy in 1976 for his work in group theory, and was awarded an honorary doctoral degree from the University of Bonn, Germany, in 1965. He had visiting appointments at the University of Chicago (1960–61), the Institute for Advanced Study at Princeton (1962–63, 1968–69), and Spring 1961), the University of Tokyo (Spring 1971), the University of Tsukuba, Osaka, and Tokyo (1984) and (1985), and the University of Padua, Italy (1994).

Walter Feit

Michio Suzuki was one of the group of brilliant young Japanese mathematicians who returned to Japan after World War II. He received his Ph.D. in 1952 from the University of Tokyo in algebra. This is that he came to the University of Illinois in 1952 as a research fellow. He joined the faculty of the University in 1953, a position he held until his sudden death.

His thesis was in the theory of finite groups, and this subject was to occupy him for the whole career. His early work included a study of the lattice of subgroups $L(G)$ of a group G . He proved that if G is a nonsolvable simple finite group and H is a finite group with $L(G \times G) = L(H \times H)$, then G is isomorphic to H . At the time it was not known whether $L(G)$ determines G up to isomorphisms. However, by using the classification of the finite simple groups, it is possible to prove the more natural result that G and H are isomorphic if and only if $L(G) = L(H)$. In 1956, then G is isomorphic to H . Consideration of a cyclic group C of prime order led to a problem of "isomorphism of lattices". He sent a letter to Feit about "isomorphism of lattices". He sent a letter to Feit about "isomorphism of lattices". He sent a letter to Feit about "isomorphism of lattices".

Photographic courtesy of Michael Aschbacher and Michio Suzuki.

Mar 1999

order indicates why $L(G \times G)$ should always be much richer than $L(G)$.

During the summer of 1952 he came to Ann Arbor, attracted by the presence of Richard Brauer, who was on the faculty there. Brauer was one of the very few senior mathematicians in the USA who worked on questions concerning the structure of finite simple groups. He and Brauer met a couple of times that summer, which John Pollard and I can still remember. I met him in that summer while I was a graduate student at the University of Michigan.

The theory of finite groups became a subject of intensive research during the next few years. One reason was John Thompson's thesis, which introduced new methods and ideas in the subject; another was the progress in character theory spearheaded by Brauer and Suzuki.

It is necessary here to make some definitions. It was of background, an important theorem that Thompson says that if H is a finite transitive permutation group such that the subgroup fixing a letter is nontrivial and no nonidentity element fixes more than one letter, then H contains a proper nontrivial normal subgroup. It says that every nonidentity element is a fixed point stabilizer contained in H . All known proofs of this theorem use character theory. We use this theorem to make a definition: A finite group H is a Frobenius group

M. Aschbacher, H. Bender, W. Feit, R. Solomon,
Michio Suzuki (1926–1998)
Notices Amer. Math. Soc. 46 (1999), no. 5, 543–551

closed abelian 2-subgroup. Thus he weakened Shult's hypotheses, but he also appealed to Shult's result in his proof. Goldschmidt's theorem is not strictly speaking a connectedness result.

Shult never published his Fusion Theorem, but one section of his proof is reproduced with slight variations in the next paper. To simplify the statement of the theorem, I assume G is simple, but this is not really necessary if one slightly extends the class of examples.

Theorem (Aschbacher [A1], [A2]). Let G be a finite simple group, H a proper subgroup of G , and $x \in H$ an involution in the center of a Sylow 2-subgroup of G . Assume

- (i) x fixes a unique point of the coset space G/H , and
- (ii) if $z \neq 1 \in x^G \cap C_G(x)$, then $C_G(xz)$ is contained in H .

Then H is strongly embedded in G , so that $G \cong L_2(2^n)$, $Sz(2^n)$, or $U_3(2^n)$.

Reading Shult's paper and Bender's paper on groups with a strongly embedded subgroup helped me to prove the result above, and that theorem is used in turn to prove the final connectedness result I will mention. Define Γ_2^3 to be the commuting graph on elementary abelian 2-subgroups of G of rank at least 2 and Γ_2^3 the subgraph of non-isolated vertices in Γ_2^3 .

Theorem (Aschbacher [A1]). Let G be a finite simple group of 2-rank at least 3 such that Γ_2^3 is disconnected. Then G is $L_3(2^n)$, $Sz(2^n)$, $U_3(2^n)$, or J_1 .

This last result is used in conjunction with signalizer functor theory to control the groups $O(C_G(x))$, Γ an involution in G . The problem of determining all such groups (phrased somewhat differently) was posed by Gorenstein and Walter. I first learned of this problem in the Suzuki-Walter seminar and treated a very special case of the problem during my year at Illinois.

Since space in this tribute is limited, I have chosen to concentrate on Suzuki's work on disconnected groups, pointing out also how he inspired his students and postdocs to take that work to its logical conclusion. Unfortunately this only hints at the many important contributions Suzuki made to the theory of finite groups. During the 1950s and 1960s Suzuki was one of the masters of the theory of group characters and one of the initiators of the local theory of finite groups. He also discovered his infinite family of simple groups, which were only later recognized to be groups of Lie type, and he discovered the sporadic Suzuki group.

I last saw Michio Suzuki in July 1997 at the conference in honor of his seventieth birthday in Tokyo. While his hair was a little grey, he was as active as ever, and it was hard to believe that he was almost thirty years older than when we first

met. Unfortunately, later that year it was discovered he was in the last stages of terminal cancer, and he died a few months after that diagnosis. His death is a great loss to mathematics and to those of us who knew him.

Helmut Bender

The name of Michio Suzuki was forever engraved in my mind when in 1964 Bernd Fischer, who had just become an assistant of Reinhold Baer at Frankfurt, handed me a paper by Suzuki to be studied and presented in Baer's seminar. That paper [Su5] lies at the intersection of two main streams of Suzuki's work:

(1) Characterize the known simple groups by the centralizer of an involution.

(2) Determine doubly transitive permutation groups with a regular fixed point behavior, especially Zassenhaus groups (a nonidentity element does not fix three points) and Suzuki-transitive groups (the stabilizer of a point has a normal subgroup regular on the remaining points).

The stream (1) was initiated by Richard Brauer in his 1954 ICM lecture and on the basis of philosophical as well as practical considerations: A given group can occur as the centralizer of an involution in only finitely many finite simple groups, up to isomorphism, and involutions allow certain arguments, elementary as well as character-theoretic ones, that do not work for elements of higher order.

Brauer's dream was that in a hypothetical future classification of the finite simple groups one would reach a point where such characterizations could be used. This dream indeed became true by Aschbacher's later work, but the relevance of Suzuki's main contributions in that area was obvious much earlier. His most general result is the determination of all finite groups in which the centralizer of any involution has a normal Sylow 2-subgroup [Su6]. The only such simple groups are $PSL_2(2^n)$, $Sz(2^n)$, and $PSL_3(2^n)$.

A further discussion of that story will carry us quickly to the main topic, 2-connectedness, of Aschbacher's segment of the present article. So, as a bridge in some other direction, let me recall a re-


Helmut Bender is professor of mathematics at the University of Kiel. His email address is bender@math.uni-kiel.de.



Michio Suzuki and daughter Kazuko at the Institute for Advanced Study, Princeton, fall of 1968.



Shiro Satake (left) and Suzuki, Japan, 1981.



Suzuki (left) with Graham Higman, July 1987.

mark Fischer once made to me in the old days, in a meeting, "Suzuki now works with BN-pairs rather than characters." This referred to (Suz) and reflects, as I understood only much later, a twofold deep inner harmony in the world of finite groups.

For a brief description of this, let me relate Suzuki's procedure in [Su6] for the classification of all finite simple groups, in either case, after some reduction of

a minimal counterexample G ,

(a) the normalizer of any nonidentity 2-subgroup (in short, any 2-local subgroup) has a normal 2-subgroup containing its centralizer, and then

(b) one either has non-2-connectedness in Aschbacher's sense or a geometry (building or BN-pair) that allows one to identify G .

In [Su6] that geometry is a projective plane with the cosets Lx ($x \in G$) of some 2-local L as lines, the cosets Fy of some other 2-local as points, incidence defined by $Lx \cap Fy \neq \emptyset$, and G acting on the plane by multiplication from the right. In either case the reduction to (a) is a big story in itself; in the general situation it involves the study of (a) also for odd primes p in place of 2, and again (b) is the goal to be reached. The way to a geometry is by the study of the p -locals and their interaction. The most exciting recent developments in finite group theory are in this field.

The main obstacle to any kind of local analysis as just indicated is non- p -connectedness; that is, some proper subgroup H of G contains a Sylow p -subgroup of G and the normalizers of all (or

nearly all nonidentity p -subgroups of H . So from the p -local point of view, H is a kind of black hole and hard to distinguish from G . As Solomon has pointed out, character theory may come to the rescue when things get very tight. Indeed, local analysis and character theory are somehow complementary methods, and it was the work of Brauer, Feit, and Suzuki on so-called exceptional characters that made this clear.

The rule of the prime 2 in the realm of finite groups is via the Sylow 2-subgroups. I like to compare them with fortresses. In order to exert an effective control, they need good lines of communication, that is, good connectedness, and most of all strength. For every type of small Sylow 2-subgroup, Suzuki has obtained substantial results. Most impressive are his CA Theorem [Su3], discussed in the segment below by Solomon,¹ and the Brauer-Suzuki-Burnside Theorem [Su1] on groups G of even order with no elementary 2-subgroup of order 4.²

After a long period of relatively little global progress, much light has been shed on finite groups in the second half of this century, not least by Suzuki and by others using his work. In particular, the finite simple groups have been determined. The earlier state of affairs was described very nicely by Brauer at the 1970 ICM: "Up to the early 1960s, really nothing of real interest was known about general simple groups of finite order." On the solvability of groups of odd order, the main cornerstone of the classification, Brauer said, "Nobody ever did anything about it, simply because nobody had any idea how to get even started."

Unfortunately, public statements about the classification are sometimes more governed by the bureaucratic mind than by mathematical interest and insight. The most exciting observation, however, is open to every mathematician, not only to experts: On the one hand, apart from sets, mappings, and integers, there is no other mathematical concept of such a general nature as the concept of a group; any mathematical structure has an automorphism group, which a priori has a good chance of being more or less simple. On the other hand, not only is there some order among the finite simple groups, but essentially all of them can be derived in a certain way from certain very special types of objects, namely, finite-dimensional Lie algebras over the complex numbers, which in their own right are of central importance for mathematics as a whole.

In today's light we see that Suzuki's work centers around the "rank 1 groups" of Lie type. Their

¹CA group is a group in which the centralizer of every nonidentity element is abelian. The CA Theorem says that CA groups of odd order are solvable.

²This theorem says that G is the product of the centralizer of an involution and a normal subgroup of odd order.

Ph.D. Students of Michio Suzuki

Steven F. Bauman (1962)
Ernest C. Shult (1964)
Anne P. Street (1966)
Jon M. Laible (1967)
John S. Montague (1967)
Robert E. Lewis (1968)
C. Gomer Thomas (1968)
Mark P. Hale (1969)
Wen-jin Wuan (1970)
Zen-I Chang (1974)
George M. Whitson (1974)
Arthur A. Yanushka (1974)
David Chillag (1975)
Seung-Ahn Park (1975)
David Redmond (1977)
Mark R. Hoekjms (1978)
Robert F. Mortenson (1978)
Michael D. Fry (1980)
Philip Abram Cobb (1987)
Randall Reed Holmes (1987)
Harald Erich Ellers (1989)
Tsung-Luen Sheu (1989)
Jose Maria Balmaceda (1991)
Tuval Foguel (1992)
Abdellatif Laradji (1993)

unifying group-theoretical property is "Suzuki-transitivity", essentially that the underlying geometry degenerates. This explains why Suzuki's main papers on permutation groups contain such long and difficult (though ingenious) calculations to identify the group under consideration. Is this effort really worthwhile? Why do group theorists always want to establish isomorphisms rather than being satisfied with the relevant group-theoretic information? This question was once raised by a mathematician in an interview. The answer is that information on some aspect of a group may suffice for an isomorphism with a known group but is usually not strong enough to clarify any other important aspect, e.g., representations. This can be illustrated by the (slightly generalized) Brauer-Suzuki-Wall Theorem [BSW]: A certain abelian subgroup of the given group G will be known to be cyclic only after an isomorphism of G with some $PSL_2(q)$ has been established.

To follow those calculations in my seminar paper was the hardest job I ever did in group theory. Fortunately, there was enough other motivation to go on and study further papers of Suzuki and related work of Feit. This became my main activity for about two years as a student. Shortly thereafter I spent a year with Suzuki at Urbana, the most productive and pleasant time in my academic life.

Once Suzuki mentioned that he came to Urbana at exactly my age. So it must have been 1952. He came on the initiative of Beer, then at Urbana, who was impressed by his work [Su1, Su2] on subgroup lattices, recently described to me by experts as revolutionary.

In 1991 Suzuki received a honorary degree from the University of Kiel for his pioneering work, more precisely "für seine Verdienste auf dem Gebiet der Gruppentheorie, vor allem in Würdigung seiner wegweisenden Arbeiten zur Klassifikation der endlichen einfachen Gruppen wie auch für sein grundlegendes Werk über Untergruppenverbände und seine Beiträge zur Theorie der Permutationsgruppen."³ The next time I saw him was in July 1997 at a meeting in Tokyo connected with his seventieth birthday. Those who know him and met him there will have no doubt that his passion for finite groups and his warm interest in the life and work of his colleagues lasted until the end of his days.

Ronald Solomon

In preparing a talk for the July 1997 conference in Tokyo in honor of Michio Suzuki, I took a moment to check the bibliography of my battered copy of Danny Gorenstein's book, *Finite Groups, (the Book of Common Prayer for students of my generation interested in finite simple groups)*. I found that the author (or coauthor) with the most citations (19) was Michio Suzuki, a crude numerical indication of the importance of Suzuki to the rapidly developing theory of finite simple groups. A far more eloquent synopsis of the influence of Michio Suzuki on the project of classifying the finite simple groups can be found in the following remarks of John Thompson:⁴


A third strategy (or was it a tactic?) in OOP [the Odd Order Paper] attempted to build a bridge from Sylow theory to character theory. The far shore was marked by the granite of Suzuki's theorem on CA-groups....

Suzuki's CA-theorem is a marvel of cunning. In order to have a genuinely satisfying proof of the odd order theo-

³For his achievements in the domain of group theory, classification of simple finite groups as well as for his fundamental work on lattices of subgroups and his contributions to the theory of permutation groups.
Ronald Solomon is professor of mathematics at Ohio State University. His e-mail address is solomon@math.ohio-state.edu.

⁴For CA groups and Suzuki's theorem that are mentioned in these remarks, see footnote 1 within Bender's segment of this article.





Photograph courtesy of Naoko Suzuki, photo copyright Naoko Suzuki

At the conference in honor of his seventieth birthday, Tokyo, 1997. Top: Suzuki with wife Naoko. Bottom: left to right, Suzuki, Donald G. Higman, and Walter Feit.

rem, it is necessary, it seems to me, not to assume this theorem. Once one accepts this theorem as a step in a general proof, one seems irresistibly drawn along the path which was followed. To my colleagues who have grumbled about the tortuous proofs in the classification of simple groups, I have a ready answer: find another proof of Suzuki's theorem.

Thus Suzuki was either Moses leading his people to the Promised Land of the classification or the Pied Piper leading a generation of thoughtless children down a tortuous path of no return.

Burstein was intrigued by the differences between groups of even and odd order and was fascinated by the thought that all groups of odd order might be solvable, but for all his brilliance he never established a major subcase of the problem. Brauer in the early 1950s focused attention on centralizers as a key to unlocking the mysteries of simple groups, but his ideas were directed primarily at centralizers of involutions and had little applicability to groups of odd order. Suzuki took up the idea of centralizer conditions (CA means that the centralizers of nonidentity elements are abelian) and

devised and implemented a strategy applicable to groups of both odd and even order:

(1) Determine the structure of all maximal local subgroups (normalizers of nonidentity p -subgroups) via Sylow theory.

(2) Count elements in G via character theory.

Suzuki's paper was seductively short and elegant, only ten pages long. Little did anyone know that it was the seed that would germinate into the 235-page Odd Order Paper of Feit and Thompson.

After the Odd Order Paper, the role of character theory in the proof of the classification declined precipitously and disappeared completely in the final decade. One heuristic explanation for this is that the primary role of character theory in the classification is to obtain group order formulas. But Thompson's Order Formula shows that the group order can be obtained in an elementary character-free way whenever the group has at least two conjugacy classes of involutions (elements of order 2). Empirically I do not know of a proof without quoting the classification; simple groups with only one class of involutions have Sylow 2-subgroups either of very small (sectional) 2-rank or of very small nilpotence class. As groups of these types were among the first handled in the classification project, it is not surprising that the need for character theory was exhausted early.

A somewhat deeper reason is the following. Character theory comes to the rescue when local group theory finds itself trapped in a cul-de-sac, a so-called strongly embedded subgroup or something very close to one. If the local structure of the group G in the neighborhood of the prime 2 is sufficiently robust to provide a family of local subgroups with large intersections which generate G , then it is possible to avoid character theory and identify G by either Lie theoretic or geometric methods. A single subgroup (and its cosets) provides a permutation action. A web of intersecting subgroups (and their cosets) provides a geometry. But the absence of rich intersections can also be exploited, as Suzuki taught us in his deepest papers. "On a class of doubly transitive groups I," here Suzuki wrote the final chapter in the long saga of Zassenhaus groups and also the first chapter of the classification of groups with a strongly embedded subgroup, which was completed by Mikhael Bender. This marvelous work complements the Odd Order Paper to give the final assurance that not only do all nonabelian finite simple groups have even order but, with the exception of the groups of Lie rank 1 over fields of even order, they are rich in 2-local subgroups. (Technically speaking, for any fixed Sylow 2-subgroup S , G is generated by the normalizers of nonidentity subgroups of S .)

This does not exhaust Suzuki's contributions to the classification. He was a mentor, official or un-

ADVANCED STUDIES
IN PURE MATHEMATICS 32

Chief Editor: Eichi Bannai (Kyushu University)

Groups and Combinatorics
— in memory of Michio Suzuki

Edited by

Eichi Bannai (Kyushu University)
Hirosaki Suzuki (International Christian University)
Hiroyoshi Yamaki (Kumamoto University) and
Tomoyuki Yoshida (Hokkaido University)

Mathematical Society of Japan

Groups and Combinatorics In Memory of Michio Suzuki

E. Bannai, H. Suzuki, H. Yamaki, T. Yoshida eds

Advanced Studies in Pure Mathematics

2001

Michio Suzuki

Koichiro Harada

51	Biographical Sketch.....	1
52	The Early Work of Michio Suzuki.....	3
53	Theory of Exceptional Characters.....	5
54	The CA-paper of Suzuki.....	7
55	Zassenhaus Groups.....	10
56	Suzuki's Simple Groups $Sz(2^n)$	13
57	2T-groups and Related Classification Theorems.....	17
58	Group Theory in Japan before Suzuki.....	29
59	Michio Suzuki, my teacher and my mentor.....	31

§1. Biographical Sketch

1926, October 2. Born in Chiba, Japan.

1942, April. Entered the Third High School of Japan located at Kyoto (Shobun In, Katsumi Nomin, Hidetshio Yamabe were his seniors by one year and Shige Marukami was in the same class).

1945, April. Entered the University of Tokyo. Majored in mathematics. (Gaisai Takeuchi, Nagayoshi Iwahori, Tetsuo Tamagawa were friends of this period.)

1948, April. Entered the Graduate School of Tokyo University. Suzuki's supervisor was Shokichi Iyanaga. Kenkichi Iwasawa had a profound influence on Suzuki.

1949-'51. Received a special graduate fellowship from the Government of Japan.

1951, April to '52, January. Held a lectureship at Tokyo University of Education.

1952, January to '52, May. Held a graduate fellowship at University of Illinois at Urbana-Champaign.

Received April 27, 1999.

Revised May 17, 1999.

Koichiro Harada



Groups and Combinatorics - In Memory of Michio Suzuki
Advanced Studies in Pure Mathematics

2001

Singapore Group Theory Conference

National University of Singapore, Singapore

June 8-19, 1987





Dr. Suzuki was a **great advisor**. He always listened to me to understand what I really had issues with and guided me to a direction in which I could comfortably excel. This was because he knew what my strengths and weaknesses are. He always had time for me and was always helpful. He installed in my the **love for group theory**.

Dr. Suzuki is best known for his work on Finite Simple groups, in fact his work was a major factor in inspiring the remarkable combined effort that led to the classification of finite simple groups which many regard as the most important mathematical achievement of the 20th century. But it is his earlier works on **partition of finite group** (1950) and on the **lattice of subgroups of finite groups** (1951) that have most influence my research to the present. I greatly appreciate his influence on my research and love of Mathematics. It was very difficult in 1998 to learn of his untimely passing just two years after my completion of my thesis under his outstanding supervision.

from **Tuval Foguel** - Adelphi University - U.S.A.

