

# Reduction of Poisson-Nijenhuis Lie algebroids

Antonio De Nicola\*

work in progress with Juan Carlos Marrero and Edith Padrón\*\*

\* CMUC, Department of Mathematics,  
University of Coimbra

\*\* Universidad de La Laguna

Poisson PT@Porto, 15 May 2010

# Overview

Aim of our study:

- Given a Poisson-Nijenhuis Lie algebroid  $(A, P, N)$  we want to reduce it to a symplectic-Nijenhuis Lie algebroid  $(\tilde{A}, \tilde{\Omega}, \tilde{N})$  with  $\tilde{\Omega}$  symplectic and also  $\tilde{N}$  nondegenerate.

# Motivation

But first a preliminary question:

- Why study Poisson-Nijenhuis Lie algebroids?

# Poisson-Nijenhuis manifolds (briefly)

**Ingredients:**  $M$  manifold,  $\Lambda$  bivector field and  $N$   $(1,1)$ -tensor on  $M$

- $\Lambda$  is a Poisson structure, i.e.,  $[\Lambda, \Lambda] = 0$
- $N$  is Nijenhuis operator, i.e.,  $\mathcal{T}_N = 0$
- $\Lambda$  and  $N$  satisfy the compatibility conditions

$$N \circ \Lambda^\sharp = \Lambda^\sharp \circ N^*, \quad C(\Lambda, N) = 0$$

where  $\Lambda^\sharp : T^*M \rightarrow TM$ ,  $\Lambda^\sharp(\alpha) = i_\alpha \Lambda$



$(M, \Lambda, N)$  Poisson-Nijenhuis manifold

# Poisson-Nijenhuis manifolds

$(M, \Lambda, N)$  Poisson-Nijenhuis manifold  $\Rightarrow \Lambda_i^\sharp = N\Lambda_{i-1}^\sharp$

Poisson structures  $\Lambda_i, \Lambda_j$  are compatible

**Particular case:**

Bi-hamiltonian manifold  $(M, \Lambda_0, \Lambda_1)$  with  $\Lambda_0$  symplectic structure



$(M, \Lambda_0, N = \Lambda_1^\sharp \circ (\Lambda_0^\sharp)^{-1})$  Poisson-Nijenhuis manifold

+

$X_1 = \Lambda_1^\sharp(dH_0) = \Lambda_0^\sharp(dH_1)$  bi-Hamiltonian vector field



$X_i = N^{i-1}X_1$  sequence of bi-Hamiltonian v. fields

A simple Example: Toda lattice (for two particles)

$\mathbb{R}^4$  with coordinates  $(q^1, q^2, p_1, p_2)$

$$H_1 = \frac{1}{2}(p_1^2 + p_2^2) + e^{q^1 - q^2}$$

Poisson structures

$$\Lambda_0 = \frac{\partial}{\partial q^1} \wedge \frac{\partial}{\partial p_1} + \frac{\partial}{\partial q^2} \wedge \frac{\partial}{\partial p_2}$$

$$\Lambda_1 = -\frac{\partial}{\partial q^1} \wedge \frac{\partial}{\partial q^2} + p_1 \frac{\partial}{\partial q^1} \wedge \frac{\partial}{\partial p_1} + p_2 \frac{\partial}{\partial q^2} \wedge \frac{\partial}{\partial p_2} + e^{q^1-q^2} \frac{\partial}{\partial p_2} \wedge \frac{\partial}{\partial p_1}$$

$$N = \Lambda_1^\# \circ (\Lambda_0^\#)^{-1}, \quad (\mathbb{R}^4, \Lambda_0, N) \text{ PN-manifold}$$

$$X_1 = \Lambda_0^\sharp(dH_1) = \Lambda_1^\sharp(dH_0), \quad H_0 = p_1 + p_2$$

# Toda lattice in Flaschka coordinates

$$\mathbb{R} \times \mathbb{R}^4 \rightarrow \mathbb{R}^4, \quad (t, (q^1, q^2, p_1, p_2)) \rightarrow (q^1 + t, q^2 + t, p_1, p_2)$$

$$\mathbb{R}^4/\mathbb{R} \cong (\mathbb{R}^+) \times \mathbb{R}^2$$

$$[(q^1, q^2, p_1, p_2)] \rightarrow (e^{q_1 - q_2}, p_1, p_2)$$

$$\pi : \mathbb{R}^4 \rightarrow \mathbb{R}^+ \times \mathbb{R}^2, \quad (q^1, q^2, p_1, p_2) \rightarrow (e^{q_1 - q_2}, p_1, p_2)$$

$(a, b_1, b_2)$  the coordinates on the reduced space  $\mathbb{R}^+ \times \mathbb{R}^2$

# Toda lattice in Flaschka coordinates

## Poisson reduced structures

$$\bar{\Lambda}_0 = a \frac{\partial}{\partial a} \wedge \left( \frac{\partial}{\partial b_1} - \frac{\partial}{\partial b_2} \right)$$

$$\bar{\Lambda}_1 = a \frac{\partial}{\partial a} \wedge \left( b_1 \frac{\partial}{\partial b_1} - b_2 \frac{\partial}{\partial b_2} \right) - a \frac{\partial}{\partial b_1} \wedge \frac{\partial}{\partial b_2}$$

$$\bar{H}_1 = \frac{1}{2}(b_1^2 + b_2^2) + a, \quad \bar{H}_0 = b_1 + b_2$$

$$\bar{X}_1 = \bar{\Lambda}_0^\sharp(d\bar{H}_1) = \bar{\Lambda}_1^\sharp(d\bar{H}_0)$$

$\exists \bar{N}$  such that  $\bar{\Lambda}_1^\sharp = \bar{N} \circ \bar{\Lambda}_0^\sharp !!!$

# What happens?

The answer is in the theory of Poisson-Nijenhuis Lie algebroids

## Lie algebroids

## Definition (Pradines)

A **Lie algebroid** is a vector bundle  $\tau_A: A \rightarrow M$  endowed with

- (i) an anchor, i.e., a vector bundle morphism  $\rho_A: A \rightarrow TM$
  - (ii) a Lie algebra bracket on  $\Gamma(A)$ ,  $[ , ]_A$ , such that

## Lie algebroids

## Definition (Pradines)

A **Lie algebroid** is a vector bundle  $\tau_A: A \rightarrow M$  endowed with

- (i) an anchor, i.e., a vector bundle morphism  $\rho_A: A \rightarrow TM$
  - (ii) a Lie algebra bracket on  $\Gamma(A)$ ,  $[ , ]_A$ , such that

$$[X, fY]_A = f [X, Y]_A + \rho_A(X)(f)Y,$$

for all  $X, Y \in \Gamma(A)$ ,  $f \in C^\infty(M)$ .

## Lie algebroids

## Definition (Pradines)

A **Lie algebroid** is a vector bundle  $\tau_A: A \rightarrow M$  endowed with

- (i) an *anchor*, i.e., a vector bundle morphism  $\rho_A: A \rightarrow TM$
  - (ii) a Lie algebra bracket on  $\Gamma(A)$ ,  $[ , ]_A$ , such that

$$[X, fY]_A = f [X, Y]_A + \rho_A(X)(f)Y,$$

for all  $X, Y \in \Gamma(A)$ ,  $f \in C^\infty(M)$ .

## Lie algebroids

## Definition (Pradines)

A **Lie algebroid** is a vector bundle  $\tau_A: A \rightarrow M$  endowed with

- (i) an *anchor*, i.e., a vector bundle morphism  $\rho_A: A \rightarrow TM$
  - (ii) a Lie algebra bracket on  $\Gamma(A)$ ,  $[ , ]_A$ , such that

$$[X, fY]_A = f [X, Y]_A + \rho_A(X)(f)Y,$$

for all  $X, Y \in \Gamma(A)$ ,  $f \in C^\infty(M)$ .

It follows that

$$\rho_A([X, Y]_A) = [\rho_A(X), \rho_A(Y)] .$$

## Examples of Lie algebroids

## The tangent bundle

$$(A = TM, \rho_A = id_{TM}, [ , ])$$

## An involutive distribution

$$(A = D \subset TM, \rho_A = \iota_D, [ , ])$$

## A Lie algebra

$$(A = \mathfrak{g}, \rho_A = 0, [\cdot, \cdot]_{\mathfrak{g}})$$

# Examples of Lie algebroids

## The Atiyah algebroid

$\pi : M \rightarrow M/G$  principal  $G$ -bundle

- $A = TM/G \rightarrow M/G$  sections are  $G$ -invariant vector fields
- $\rho_A([v]) = T\pi(v)$  induced projection map
- $[ , ]_A$  = bracket of  $G$ -invariant vector fields

# Cartan calculus

Associated to a given Lie algebroid  $(A, [\cdot, \cdot]_A, \rho_A)$  there is a *Lie algebroid differential*  $d^A: \Gamma(\wedge^\bullet A^*) \rightarrow \Gamma(\wedge^{\bullet+1} A^*)$  defined by

$$(d^A\omega)(X_0, \dots, X_k) := \sum_{i=0}^k (-1)^i \rho_A(X_i) \left( \omega(X_0, \dots, \hat{X}_i, \dots, X_k) \right) + \sum_{0 \leq i < j \leq k} (-1)^{i+j} \omega([X_i, X_j]_A, X_0, \dots, \hat{X}_i, \dots, \hat{X}_j, \dots, X_k),$$

for  $\omega \in \Gamma(\wedge^k A^*)$ ,  $X_0, \dots, X_k \in \Gamma(A)$ .

- For  $X \in \Gamma(A)$ ,

$$\mathcal{L}_X^A := i_X d^A + d^A i_X$$

# Cartan calculus

Associated to a given Lie algebroid  $(A, [\cdot, \cdot]_A, \rho_A)$  there is a *Lie algebroid differential*  $d^A: \Gamma(\wedge^\bullet A^*) \rightarrow \Gamma(\wedge^{\bullet+1} A^*)$  defined by

$$(d^A\omega)(X_0, \dots, X_k) := \sum_{i=0}^k (-1)^i \rho_A(X_i) \left( \omega(X_0, \dots, \hat{X}_i, \dots, X_k) \right) + \sum_{0 \leq i < j \leq k} (-1)^{i+j} \omega([X_i, X_j]_A, X_0, \dots, \hat{X}_i, \dots, \hat{X}_j, \dots, X_k),$$

for  $\omega \in \Gamma(\wedge^k A^*)$ ,  $X_0, \dots, X_k \in \Gamma(A)$ .

- For  $X \in \Gamma(A)$ ,

$$\mathcal{L}_X^A := i_X d^A + d^A i_X$$

# Properties of the Lie algebroid differential

- $d^A$  is a graded derivation of degree 1, i.e.,

$$d^A(\theta \wedge \omega) = d^A\theta \wedge \omega + (-1)^{\deg(\theta)}\theta \wedge d^A\omega,$$

- $d^A \circ d^A = 0.$

# Properties of the Lie algebroid differential

- $d^A$  is a graded derivation of degree 1, i.e.,

$$d^A(\theta \wedge \omega) = d^A\theta \wedge \omega + (-1)^{\deg(\theta)}\theta \wedge d^A\omega,$$

- $d^A \circ d^A = 0$ .

# Morphisms of Lie algebroids

Let  $(A, [\cdot, \cdot]_A, \rho_A)$  and  $(A', [\cdot, \cdot]_{A'}, \rho_{A'})$  be Lie algebroids. A bundle map

$$\begin{array}{ccc} A & \xrightarrow{F} & A' \\ \tau_A \downarrow & & \downarrow \tau_{A'} \\ M & \xrightarrow{f} & M' \end{array}$$

is called a **morphism of Lie algebroids** from  $A$  to  $A'$ , if

$$d^A(F^*\alpha') = F^*(d^{A'}\alpha') \quad \text{for all } \alpha' \in \Gamma(\wedge^k A'^*).$$

## Schouten-Gerstenhaber algebra

The Lie algebra bracket on  $\Gamma(A)$  can be extended to the exterior algebra  $(\Gamma(\wedge^\bullet A), \wedge)$ . For  $X \in \Gamma(A)$  and  $P \in \Gamma(\wedge^p A)$ ,

$$[X, P]_A(\alpha_1, \dots, \alpha_p) = \rho_A(X)(P(\alpha_1, \dots, \alpha_p))$$

$$- \sum_{i=1}^p P(\alpha_1, \dots, \mathcal{L}_X^A \alpha_i, \dots \alpha_p),$$

## Schouten-Gerstenhaber algebra

The Lie algebra bracket on  $\Gamma(A)$  can be extended to the exterior algebra  $(\Gamma(\wedge^\bullet A), \wedge)$ . For  $X \in \Gamma(A)$  and  $P \in \Gamma(\wedge^p A)$ ,

$$[X, P]_A(\alpha_1, \dots, \alpha_p) = \rho_A(X)(P(\alpha_1, \dots, \alpha_p))$$

$$- \sum_{i=1}^p P(\alpha_1, \dots, \mathcal{L}_X^A \alpha_i, \dots \alpha_p),$$

If  $P \in \Gamma(\wedge^p A)$ ,  $Q \in \Gamma(\wedge^q A)$  and  $R \in \Gamma(\wedge^r A)$ , then  
 $[P, Q]_A \in \Gamma(\wedge^{p+q-1} A)$  and

- $[P, Q]_A = -(-1)^{(p-1)(q-1)} [Q, P]_A$
  - $[P, Q \wedge R]_A = [P, Q]_A \wedge R + (-1)^{(p-1)q} Q \wedge [P, R]_A$
  - $(-1)^{(p-1)(r-1)} [P, [Q, R]_A]_A + \text{cyclic perm.} = 0$

## Schouten-Gerstenhaber algebra

The Lie algebra bracket on  $\Gamma(A)$  can be extended to the exterior algebra  $(\Gamma(\wedge^\bullet A), \wedge)$ . For  $X \in \Gamma(A)$  and  $P \in \Gamma(\wedge^p A)$ ,

$$[X, P]_A(\alpha_1, \dots, \alpha_p) = \rho_A(X)(P(\alpha_1, \dots, \alpha_p))$$

$$- \sum_{i=1}^p P(\alpha_1, \dots, \mathcal{L}_X^A \alpha_i, \dots \alpha_p),$$

If  $P \in \Gamma(\wedge^p A)$ ,  $Q \in \Gamma(\wedge^q A)$  and  $R \in \Gamma(\wedge^r A)$ , then  
 $[P, Q]_A \in \Gamma(\wedge^{p+q-1} A)$  and

- $[P, Q]_A = -(-1)^{(p-1)(q-1)} [Q, P]_A$
  - $[P, Q \wedge R]_A = [P, Q]_A \wedge R + (-1)^{(p-1)q} Q \wedge [P, R]_A$
  - $(-1)^{(p-1)(r-1)} [P, [Q, R]_A]_A + \text{cyclic perm.} = 0$

# Schouten-Gerstenhaber algebra

The Lie algebra bracket on  $\Gamma(A)$  can be extended to the exterior algebra  $(\Gamma(\wedge^\bullet A), \wedge)$ . For  $X \in \Gamma(A)$  and  $P \in \Gamma(\wedge^p A)$ ,

$$[X, P]_A(\alpha_1, \dots, \alpha_p) = \rho_A(X)(P(\alpha_1, \dots, \alpha_p)) - \sum_{i=1}^p P(\alpha_1, \dots, \mathcal{L}_X^A \alpha_i, \dots, \alpha_p),$$

If  $P \in \Gamma(\wedge^p A)$ ,  $Q \in \Gamma(\wedge^q A)$  and  $R \in \Gamma(\wedge^r A)$ , then  $[P, Q]_A \in \Gamma(\wedge^{p+q-1} A)$  and

- $[P, Q]_A = -(-1)^{(p-1)(q-1)} [Q, P]_A$
- $[P, Q \wedge R]_A = [P, Q]_A \wedge R + (-1)^{(p-1)q} Q \wedge [P, R]_A$
- $(-1)^{(p-1)(r-1)} [P, [Q, R]_A]_A + \text{cyclic perm.} = 0$

# Poisson structures on Lie algebroids

Let  $A$  be a Lie algebroid and  $P$  a section of the vector bundle  $\wedge^2 A \rightarrow M$ . We denote by  $P^\sharp$  the usual bundle map

$$P^\sharp: A^* \longrightarrow A: \alpha \longmapsto P^\sharp(\alpha) = i_\alpha P.$$

## Definition

A **Poisson structure** on  $A$  is a section  $P \in \Gamma(\wedge^2 A)$ , such that

$$[P, P]_A = 0.$$

In this case, the bracket

$$[\alpha, \beta]_P := \mathcal{L}_{P^\sharp \alpha}^A \beta - \mathcal{L}_{P^\sharp \beta}^A \alpha - d^A(P(\alpha, \beta)), \quad \alpha, \beta \in \Gamma(A^*),$$

is a Lie bracket and  $A_P^* = (A^*, [\cdot, \cdot]_P, \rho_A \circ P^\sharp)$  is a Lie algebroid.

# Poisson structures on Lie algebroids

Let  $A$  be a Lie algebroid and  $P$  a section of the vector bundle  $\wedge^2 A \rightarrow M$ . We denote by  $P^\sharp$  the usual bundle map

$$P^\sharp: A^* \longrightarrow A: \alpha \longmapsto P^\sharp(\alpha) = i_\alpha P.$$

## Definition

A **Poisson structure** on  $A$  is a section  $P \in \Gamma(\wedge^2 A)$ , such that

$$[P, P]_A = 0.$$

In this case, the bracket

$$[\alpha, \beta]_P := \mathcal{L}_{P^\sharp \alpha}^A \beta - \mathcal{L}_{P^\sharp \beta}^A \alpha - d^A(P(\alpha, \beta)), \quad \alpha, \beta \in \Gamma(A^*),$$

is a Lie bracket and  $A_P^* = (A^*, [\cdot, \cdot]_P, \rho_A \circ P^\sharp)$  is a Lie algebroid.

# Nijenhuis operators

Let  $(A, [\cdot, \cdot], \rho_A)$  be a Lie algebroid and  $N : A \rightarrow A$  a bundle map.  
The torsion of  $N$  is defined by

$$\mathcal{T}_N(X, Y) := [NX, NY]_A - N[X, Y]_A, \quad X, Y \in \Gamma(A),$$

where

$$[X, Y]_N := [NX, Y]_A + [X, NY]_A - N[X, Y]_A, \quad X, Y \in \Gamma(A).$$

When  $\mathcal{T}_N = 0$ ,  $N$  is called a *Nijenhuis operator*,  
 $A_N = (A, [\cdot, \cdot]_N, \rho_N = \rho_A \circ N)$  is a new Lie algebroid and

$$N : A_N \rightarrow A$$

is a Lie algebroid morphism.

# Nijenhuis operators

Let  $(A, [\cdot, \cdot], \rho_A)$  be a Lie algebroid and  $N : A \rightarrow A$  a bundle map. The torsion of  $N$  is defined by

$$\mathcal{T}_N(X, Y) := [NX, NY]_A - N[X, Y]_N, \quad X, Y \in \Gamma(A),$$

where

$$[X, Y]_N := [NX, Y]_A + [X, NY]_A - N[X, Y]_A, \quad X, Y \in \Gamma(A).$$

When  $\mathcal{T}_N = 0$ ,  $N$  is called a **Nijenhuis operator**,  
 $A_N = (A, [\cdot, \cdot]_N, \rho_N = \rho_A \circ N)$  is a new Lie algebroid and

$$N : A_N \rightarrow A$$

is a Lie algebroid morphism.

# Poisson-Nijenhuis Lie algebroids

On a Lie algebroid  $A$  with a Poisson structure  $P \in \Gamma(\wedge^2 A)$ , we say that a bundle map  $N : A \rightarrow A$  is **compatible** with  $P$  if

- (i)  $NP^\sharp = P^\sharp N^*$ ,
- (ii)  $[\alpha, \beta]_{NP} - [\alpha, \beta]_P^{N^*} = 0$ ,

## Definition (Grabowski-Urbanski)

A **Poisson-Nijenhuis Lie algebroid**  $(A, P, N)$  is a Lie algebroid  $A$  equipped with a Poisson structure  $P$  and a Nijenhuis operator  $N : A \rightarrow A$  compatible with  $P$ .

# Poisson-Nijenhuis Lie algebroids

On a Lie algebroid  $A$  with a Poisson structure  $P \in \Gamma(\wedge^2 A)$ , we say that a bundle map  $N : A \rightarrow A$  is **compatible** with  $P$  if

- (i)  $NP^\sharp = P^\sharp N^*$ ,
- (ii)  $[\alpha, \beta]_{NP} - [\alpha, \beta]_P^{N^*} = 0$ ,

where  $[\cdot, \cdot]_{NP}$  is the bracket defined by  $NP \in \Gamma(\wedge^2 A)$ , and  $[\cdot, \cdot]_P^{N^*}$  is the bracket obtained from  $[\cdot, \cdot]_P$  by deformation along  $N^*$ .

## Definition (Grabowski-Urbanski)

A **Poisson-Nijenhuis Lie algebroid**  $(A, P, N)$  is a Lie algebroid  $A$  equipped with a Poisson structure  $P$  and a Nijenhuis operator  $N : A \rightarrow A$  compatible with  $P$ .

# Poisson-Nijenhuis Lie algebroids

On a Lie algebroid  $A$  with a Poisson structure  $P \in \Gamma(\wedge^2 A)$ , we say that a bundle map  $N : A \rightarrow A$  is **compatible** with  $P$  if

- (i)  $NP^\sharp = P^\sharp N^*$ ,
- (ii)  $[\alpha, \beta]_{NP} - [\alpha, \beta]_P^{N^*} = 0$ ,

where  $[\cdot, \cdot]_{NP}$  is the bracket defined by  $NP \in \Gamma(\wedge^2 A)$ , and  $[\cdot, \cdot]_P^{N^*}$  is the bracket obtained from  $[\cdot, \cdot]_P$  by deformation along  $N^*$ .

## Definition (Grabowski-Urbanski)

A **Poisson-Nijenhuis Lie algebroid**  $(A, P, N)$  is a Lie algebroid  $A$  equipped with a Poisson structure  $P$  and a Nijenhuis operator  $N : A \rightarrow A$  compatible with  $P$ .

# Poisson-Nijenhuis Lie algebroids

On a Lie algebroid  $A$  with a Poisson structure  $P \in \Gamma(\wedge^2 A)$ , we say that a bundle map  $N : A \rightarrow A$  is **compatible** with  $P$  if

- (i)  $NP^\sharp = P^\sharp N^*$ ,
- (ii)  $[\alpha, \beta]_{NP} - [\alpha, \beta]_P^{N^*} = 0$ ,

where  $[\cdot, \cdot]_{NP}$  is the bracket defined by  $NP \in \Gamma(\wedge^2 A)$ , and  $[\cdot, \cdot]_P^{N^*}$  is the bracket obtained from  $[\cdot, \cdot]_P$  by deformation along  $N^*$ .

## Definition (Grabowski-Urbanski)

A **Poisson-Nijenhuis Lie algebroid**  $(A, P, N)$  is a Lie algebroid  $A$  equipped with a Poisson structure  $P$  and a Nijenhuis operator  $N : A \rightarrow A$  compatible with  $P$ .

# Toda lattice in Flaschka coordinates

$\mathbb{R}^+ \times \mathbb{R}^2$  with coordinates  $(a, b_1, b_2)$

$$\pi : \mathbb{R}^4 \rightarrow \mathbb{R}^+ \times \mathbb{R}^2$$

## Poisson reduced structures on $\mathbb{R}^+ \times \mathbb{R}^2$

$$\bar{\Lambda}_0 = a \frac{\partial}{\partial a} \wedge \left( \frac{\partial}{\partial b_1} - \frac{\partial}{\partial b_2} \right)$$

$$\bar{\Lambda}_1 = a \frac{\partial}{\partial a} \wedge \left( b_1 \frac{\partial}{\partial b_1} - b_2 \frac{\partial}{\partial b_2} \right) - a \frac{\partial}{\partial b_1} \wedge \frac{\partial}{\partial b_2}$$

## Toda lattice in Flaschka coordinates

$$T(\mathbb{R}^+ \times \mathbb{R}^2) \rightarrow \mathbb{R}^+ \times \mathbb{R}^2, \quad ([\cdot, \cdot], Id)$$

## The Lie algebroid

$$A = \mathbb{R} \times T(\mathbb{R}^+ \times \mathbb{R}^2) \rightarrow \mathbb{R}^+ \times \mathbb{R}^2$$

$$\{e_0 = (1, 0), e_1 = \left(0, \frac{\partial}{\partial a}\right), e_2 = \left(0, \frac{\partial}{\partial b_1}\right), e_3 = \left(0, \frac{\partial}{\partial b_2}\right)\}$$

$$[e_i, e_j]_A = 0, \quad \rho(e_0) = 0, \quad \rho(e_1) = \frac{\partial}{\partial a}, \quad \rho(e_2) = \frac{\partial}{\partial b_1}, \quad \rho(e_3) = \frac{\partial}{\partial b_2}$$

## Two Poisson structures on the Lie algebroid $A = \mathbb{R} \times T(\mathbb{R}^+ \times \mathbb{R}^2)$

$$P_0 = ae_1 \wedge (e_2 - e_3) + e_0 \wedge e_3$$

$$P_1 = ae_0 \wedge e_1 + ae_1(b_1e_2 - b_2e_3) + ae_2 \wedge e_3 + b_2e_0 \wedge e_3$$

These Poisson structures induce  $\bar{\Lambda}_0, \bar{\Lambda}_1$  on  $\mathbb{R}^+ \times \mathbb{R}^2$ .

## Toda lattice in Flaschka coordinates

The Nijenhuis operator  $N$

$$N = P_1^\sharp \circ (P_0^\sharp)^{-1} : A \rightarrow A$$

↓

The Poisson-Nijenhuis Lie algebroid

$$(A = \mathbb{R} \times T(\mathbb{R}^+ \times \mathbb{R}^2) \rightarrow \mathbb{R}^+ \times \mathbb{R}^2, P_0, N)$$

$$P_0^\# d^A \bar{H}_1 = P_1^\# d^A \bar{H}_0$$

$$\bar{\Lambda}_0^\sharp d\bar{H}_1 = \rho_A(P_0^\sharp d^A \bar{H}_1) = \rho_A(P_1^\sharp d^A \bar{H}_0) = \bar{\Lambda}_1^\sharp d\bar{H}_0$$

R. Caseiro: Modular classes of Poisson-Nijenhuis Lie algebroids, Lett. Math. Phys. **80** (2007) 223–238

Poisson-Nijenhuis Lie algebroids

## General case

$\pi : M \rightarrow M/G$  principal bundle

### ( $M, P, N$ ) PN-manifold

P-N G-invariants

$$\tilde{\pi} : TM/G \rightarrow M/G$$

## Atiyah algebroid

(P, N) PN-Lie algebroid

$M/G$  is not, in general,  
PN-manifold!!

Toda lattice

$\pi : \mathbb{R}^4 \rightarrow \mathbb{R}^4/\mathbb{R}$  is  $\mathbb{R}^+ \times \mathbb{R}^2$  principal bundle

$(\mathbb{R}^4, P, N)$  PN-manifold

P-N  $\mathbb{R}$ -invariants

$$\tilde{\pi} : T\mathbb{R}^4/\mathbb{R} \cong \mathbb{R} \times T(\mathbb{R}^+ \times \mathbb{R}^2) \rightarrow \mathbb{R}^+ \times \mathbb{R}^2$$

## Atiyah algebroid

(P, N) PN-Lie algebroid

$\mathbb{R}^+ \times \mathbb{R}^2$  is not PN-manifold!!

# Reduction of PN Lie algebroids

$(A, P, N)$

Poisson-Nijenhuis Lie algebroid

$\Downarrow$  Reduction by restriction

$(\bar{A}, \bar{P}, \bar{N})$

symplectic-Nijenhuis Lie algebroid

$\Downarrow$  Reduction by projection

$(\tilde{A}, \tilde{P}, \tilde{N})$

symplectic-Nijenhuis Lie algebroid with  $\tilde{N}$  nondegenerate

i.e.  $\tilde{P}^\sharp : \tilde{A}^* \rightarrow \tilde{A}$ ,  $\tilde{N} : \tilde{A} \rightarrow \tilde{A}$  isomorphisms

## 1<sup>st</sup> step: Reduction by restriction

$(A, [\cdot, \cdot]_A, \rho_A, P, N)$  Poisson-Nijenhuis Lie algebroid on  $M$ .

$D$  is locally finitely generated

## 1<sup>st</sup> step: Reduction by restriction

$(A, [\cdot, \cdot]_A, \rho_A, P, N)$  Poisson-Nijenhuis Lie algebroid on  $M$ .

Distribution  $D \subset TM$ ,  $D(x) := \rho_A(P^\sharp(A_x^*)) \subset T_x M$  for  $x \in M$

$D$  is locally finitely generated

## 1<sup>st</sup> step: Reduction by restriction

$(A, [\cdot, \cdot]_A, \rho_A, P, N)$  Poisson-Nijenhuis Lie algebroid on  $M$ .

Distribution  $D \subset TM$ ,  $D(x) := \rho_A(P^\sharp(A_x^*)) \subset T_x M$  for  $x \in M$

$$\left[ \rho_A(P^\sharp\alpha), \rho_A(P^\sharp\beta) \right] = \rho_A(P^\sharp [\alpha, \beta]_P)$$

$D$  is locally finitely generated

## 1<sup>st</sup> step: Reduction by restriction

$(A, [\cdot, \cdot]_A, \rho_A, P, N)$  Poisson-Nijenhuis Lie algebroid on  $M$ .

Distribution  $D \subset TM$ ,  $D(x) := \rho_A(P^\sharp(A_x^*)) \subset T_x M$  for  $x \in M$

$$\left[ \rho_A(P^\sharp\alpha), \rho_A(P^\sharp\beta) \right] = \rho_A(P^\sharp [\alpha, \beta]_P)$$

$D$  is locally finitely generated



$D$  is a generalized foliation of  $M$  in the sense of Sussmann.

## 1<sup>st</sup> step: Reduction by restriction

- Let  $L \subset M$  be a leaf of the foliation  $D = \rho_A(P^\sharp(A^*)) \subset TM$
  - Assume:  $P^\sharp : A^* \rightarrow A$  has constant rank on each leaf  $L$ .

↓

$A_L := P^\sharp(A^*)|_L \subset A \rightarrow L$  is a Lie algebroid

- $[P^\sharp \alpha|_L, P^\sharp \beta|_L]_{A_L} = P^\sharp [\alpha, \beta]_{P|L} \in \Gamma(A_L)$
  - $\rho_{A_L} = (\rho_A)|_{A_L} : A_L \rightarrow TL$

## 1<sup>st</sup> step: Reduction by restriction

Furthermore, the inclusion maps

$$\begin{array}{ccc} A_L & \xrightarrow{\quad I \quad} & A \\ \downarrow (\tau_A)_{|A_L} & & \downarrow \tau_A \\ L & \xrightarrow{\quad \iota \quad} & M \end{array}$$

give a morphism of Lie algebroids, i.e.

$A_L \rightarrow L$  is a Lie subalgebroid of  $A \rightarrow M$ .

# 1<sup>st</sup> step: Reduction by restriction

$$\begin{array}{ccc} L & \xrightarrow{X_L} & A_L \\ \downarrow \iota & & \downarrow I \\ M & \xrightarrow{P^\sharp \alpha} & A \end{array}$$

$$\alpha \in \Gamma(A^*)$$

The symplectic structure  $\Omega_L : L \rightarrow \wedge^2 A_L^*$

$$\Omega(X_L, Y_L) = P(\alpha, \beta) \circ \iota$$

Nijenhuis tensor  $N_L : A_L \rightarrow A_L$

$$I \circ N_L(X_L) = N(P^\sharp \alpha) \circ \iota$$

# 1<sup>st</sup> step: Reduction by restriction

## Theorem 1

Let  $(A, P, N)$  be a Poisson-Nijenhuis Lie algebroid such that the Poisson structure has constant rank in the leaves of the foliation  $D = \rho_A(P^\sharp(A^*))$ . Then, we have a symplectic-Nijenhuis Lie algebroid  $(A_L, \Omega_L, N_L)$  on each leaf  $L$  of  $D$ .

## 2<sup>nd</sup> step: Reduction by projection

### Lie algebroid epimorphism

Let  $\tau_A: A \rightarrow M$  and  $\tau_{\tilde{A}}: \tilde{A} \rightarrow \tilde{M}$  be Lie algebroids

$$\begin{array}{ccc} A & \xrightarrow{\Pi} & \tilde{A} \\ \downarrow \tau_A & & \downarrow \tau_{\tilde{A}} \\ M & \xrightarrow{\pi} & \tilde{M} \end{array}$$

- $(\Pi, \pi)$  epimorphism of vector bundles
- $d^A(\Pi^*\tilde{\alpha}) = \Pi^*(d^{\tilde{A}}\tilde{\alpha})$  for all  $\tilde{\alpha} \in \Gamma(\wedge^k \tilde{A}^*)$  and all  $k$

# Projectability

$$\begin{array}{ccc}
 A & \xrightarrow{\Pi} & \tilde{A} \\
 \downarrow \tau_A & & \downarrow \tau_{\tilde{A}} \\
 M & \xrightarrow{\pi} & \tilde{M}
 \end{array}$$

$\swarrow X \quad \nearrow \tilde{X}$

- **$\Pi$ -projectable 1-section:**  $X \in \Gamma(A)$  such that there exists  $\tilde{X} \in \Gamma(\tilde{A})$  with  $\Pi \circ X = \tilde{X} \circ \pi$ .
- **$\Pi$ -projectable 2-section:**  $P \in \Gamma(\wedge^2 A)$  such that for all  $\tilde{\alpha} \in \Gamma(\tilde{A}^*)$  the 1-section  $P^\#(\Pi^* \tilde{\alpha}) \in \Gamma(A)$  is  $\Pi$ -projectable

↓

$$\tilde{P} \in \Gamma(\wedge^2 \tilde{A}), \quad (\tilde{P}^\# \tilde{\alpha}) \circ \pi = \Pi(P^\#(\Pi^* \tilde{\alpha})).$$

- **$\Pi$ -projectable  $(1,1)$ -section:**  $N: A \rightarrow A$  vector bundle morphism such that

$$N(\Gamma_p(A)) \subseteq \Gamma_p(A) \quad \text{and} \quad \downarrow N(\Gamma(\text{Ker}\Pi)) \subseteq \Gamma(\text{Ker}\Pi)$$

$$\tilde{N}: \tilde{A} \rightarrow \tilde{A}, \quad (\tilde{N}\tilde{X}) \circ \pi = \Pi(NX).$$

# Projectability

$$\begin{array}{ccc}
 A & \xrightarrow{\Pi} & \tilde{A} \\
 \downarrow \pi & & \downarrow \tau_{\tilde{A}} \\
 M & \xrightarrow{\pi} & \tilde{M}
 \end{array}$$

- **$\Pi$ -projectable 1-section:**  $X \in \Gamma(A)$  such that there exists  $\tilde{X} \in \Gamma(\tilde{A})$  with  $\Pi \circ X = \tilde{X} \circ \pi$ .
- **$\Pi$ -projectable 2-section:**  $P \in \Gamma(\wedge^2 A)$  such that for all  $\tilde{\alpha} \in \Gamma(\tilde{A}^*)$  the 1-section  $P^\sharp(\Pi^* \tilde{\alpha}) \in \Gamma(A)$  is  $\Pi$ -projectable

$\Downarrow$

$$\tilde{P} \in \Gamma(\wedge^2 \tilde{A}), \quad (\tilde{P}^\sharp \tilde{\alpha}) \circ \pi = \Pi(P^\sharp(\Pi^* \tilde{\alpha})).$$

- **$\Pi$ -projectable  $(1,1)$ -section:**  $N: A \rightarrow A$  vector bundle morphism such that

$$N(\Gamma_p(A)) \subseteq \Gamma_p(A) \quad \text{and} \quad \downarrow N(\Gamma(\text{Ker}\Pi)) \subseteq \Gamma(\text{Ker}\Pi)$$

$$\tilde{N}: \tilde{A} \rightarrow \tilde{A}, \quad (\tilde{N} \tilde{X}) \circ \pi = \Pi(NX).$$

## Projectability

$$\begin{array}{ccc} A & \xrightarrow{\Pi} & \tilde{A} \\ \downarrow \tau_A & & \downarrow \tau_{\tilde{A}} \\ M & \xrightarrow{\pi} & \tilde{M} \end{array}$$

- **$\Pi$ -projectable 1-section:**  $X \in \Gamma(A)$  such that there exists  $\tilde{X} \in \Gamma(\tilde{A})$  with  $\Pi \circ X = \tilde{X} \circ \pi$ .
  - **$\Pi$ -projectable 2-section:**  $P \in \Gamma(\wedge^2 A)$  such that for all  $\tilde{\alpha} \in \Gamma(\tilde{A}^*)$  the 1-section  $P^\#(\Pi^* \tilde{\alpha}) \in \Gamma(A)$  is  $\Pi$ -projectable

$$\tilde{P} \in \Gamma(\wedge^2 \tilde{A}), \quad (\tilde{P}^\# \tilde{\alpha}) \circ \pi = \Pi(P^\#(\Pi^* \tilde{\alpha})).$$

- **$\Pi$ -projectable  $(1, 1)$ -section:**  $N: A \rightarrow A$  a vector bundle morphism such that

$$N(\Gamma_p(A)) \subseteq \Gamma_p(A) \quad \text{and} \quad \downarrow N(\Gamma(Ker\Pi)) \subseteq \Gamma(Ker\Pi)$$

$$\widetilde{N} : \widetilde{A} \rightarrow \widetilde{A}, \quad (\widetilde{N}\widetilde{X}) \circ \pi = \Pi(NX).$$

# Reduction by epimorphisms of Lie algebroids

$$\begin{array}{ccc} A & \xrightarrow{\Pi} & \tilde{A} \\ \downarrow \tau_A & & \downarrow \tau_{\tilde{A}} \\ M & \xrightarrow{\pi} & \tilde{M} \end{array}$$

## Theorem

Let  $(\Pi, \pi) : A \rightarrow \tilde{A}$  be a Lie algebroid epimorphism. Assume that  $(P, N)$  is a Poisson-Nijenhuis structure on  $A$  such that  $P$  and  $N$  are  $\Pi$ -projectable. Then,  $(\tilde{P}, \tilde{N})$  is a Poisson-Nijenhuis structure on  $\tilde{A}$ .

# Complete and vertical lifts

- $(A, [\cdot, \cdot]_A, \rho_A)$  a Lie algebroid
- $X \in \Gamma(A)$

The vertical lift of  $X$ :  $X^\nu \in \mathfrak{X}(A)$

- (i)  $X^\nu(f \circ \tau_A) = 0, \quad f \in \mathcal{C}^\infty(M),$
- (ii)  $X^\nu(\hat{\alpha}) = \alpha(X) \circ \tau_A, \quad \alpha \in \Gamma(A^*).$

Here, if  $\alpha \in \Gamma(A^*)$  then  $\hat{\alpha}: A \rightarrow \mathbb{R}$  is defined by

$$\hat{\alpha}(a) = \alpha(\tau_A(a))(a), \quad \text{for all } a \in A.$$

# Complete and vertical lifts

- $(A, [\cdot, \cdot]_A, \rho_A)$  a Lie algebroid
- $X \in \Gamma(A)$

The vertical lift of  $X$ :  $X^v \in \mathfrak{X}(A)$

- (i)  $X^v(f \circ \tau_A) = 0, \quad f \in \mathcal{C}^\infty(M),$
- (ii)  $X^v(\hat{\alpha}) = \alpha(X) \circ \tau_A, \quad \alpha \in \Gamma(A^*).$

The complete lift of  $X$ :  $X^c \in \mathfrak{X}(A)$

- (i)  $X^c(f \circ \tau_A) = \rho_A(X)(f) \circ \tau_A, \quad f \in \mathcal{C}^\infty(M),$
- (ii)  $X^c(\hat{\alpha}) = \widehat{\mathcal{L}_X^A} \alpha, \quad \alpha \in \Gamma(A^*).$

Here, if  $\alpha \in \Gamma(A^*)$  then  $\hat{\alpha}: A \rightarrow \mathbb{R}$  is defined by

$$\hat{\alpha}(a) = \alpha(\tau_A(a))(a), \quad \text{for all } a \in A.$$

Reduction by lifts of sections of a Lie subalgebroid

Let  $\tau_A: A \rightarrow M$  a vector bundle and  $(A, [\cdot, \cdot]_A, \rho_A)$  a Lie algebroid. Consider a Lie subalgebroid  $\tau_B: B \rightarrow M$  of  $A$ .

## Key Fact

The distributions  $\rho_A(B)$  and  $\mathcal{F}$  defined by

$$\mathcal{F}_a := \{X^c(a) + Y^\nu(a) \mid X, Y \in \Gamma(B)\} \subseteq T_a A, \quad \text{for all } a \in A.$$

are generalized foliations.

Now assume that

Reduction by lifts of sections of a Lie subalgebroid

Let  $\tau_A: A \rightarrow M$  a vector bundle and  $(A, [\cdot, \cdot]_A, \rho_A)$  a Lie algebroid. Consider a Lie subalgebroid  $\tau_B: B \rightarrow M$  of  $A$ .

## Key Fact

The distributions  $\rho_A(B)$  and  $\mathcal{F}$  defined by

$$\mathcal{F}_a := \{X^c(a) + Y^\nu(a) \mid X, Y \in \Gamma(B)\} \subseteq T_a A, \quad \text{for all } a \in A$$

are generalized foliations.

Now assume that

- (i)  $\rho_A(B)$  and  $\mathcal{F}$  are regular foliations;
  - (ii) For all  $x \in M$ ,  $a_x, a'_x \in L_{\mathcal{F}} \implies a_x - a'_x \in B_x$ .

# Reduction by lifts of sections of a Lie subalgebroid

Let  $\tau_A: A \rightarrow M$  a vector bundle and  $(A, [\cdot, \cdot]_A, \rho_A)$  a Lie algebroid.  
Consider a Lie subalgebroid  $\tau_B: B \rightarrow M$  of  $A$ .

## Key Fact

The distributions  $\rho_A(B)$  and  $\mathcal{F}$  defined by

$$\mathcal{F}_a := \{X^c(a) + Y^\nu(a) \mid X, Y \in \Gamma(B)\} \subseteq T_a A, \quad \text{for all } a \in A$$

are generalized foliations.

Now assume that

- (i)  $\rho_A(B)$  and  $\mathcal{F}$  are regular foliations;
- (ii) For all  $x \in M$ ,  $a_x, a'_x \in L_{\mathcal{F}} \implies a_x - a'_x \in B_x$ .

# Reduction by lifts of sections of a Lie subalgebroid

We define  $\tau_{\tilde{A}}: \tilde{A} = A/\mathcal{F} \rightarrow \tilde{M} = M/\rho_A(B)$  such that the following diagram is commutative

$$\begin{array}{ccc} A & \xrightarrow{\Pi} & \tilde{A} = A/\mathcal{F} \\ \downarrow \tau_A & & \downarrow \tau_{\tilde{A}} \\ M & \xrightarrow{\pi} & \tilde{M} = M/\rho_A(B) \end{array}$$

## Proposition

*In the above conditions we can define a Lie algebroid structure on*

$$\tau_{\tilde{A}}: \tilde{A} = A/\mathcal{F} \rightarrow \tilde{M} = M/\rho_A(B)$$

*such that the above diagram is an epimorphism of Lie algebroids.*

## Reduction by lifts of sections of a Lie subalgebroid

We define  $\tau_{\tilde{A}}: \tilde{A} = A/\mathcal{F} \rightarrow \tilde{M} = M/\rho_A(B)$  such that the following diagram is commutative

$$\begin{array}{ccc} A & \xrightarrow{\Pi} & \tilde{A} = A/\mathcal{F} \\ \downarrow \tau_A & & \downarrow \tau_{\tilde{A}} \\ M & \xrightarrow{\pi} & \tilde{M} = M/\rho_A(B) \end{array}$$

### Proposition

*In the above conditions we can define a Lie algebroid structure on*

$$\tau_{\tilde{A}}: \tilde{A} = A/\mathcal{F} \rightarrow \tilde{M} = M/\rho_A(B)$$

*such that the above diagram is an epimorphism of Lie algebroids.*

# The Riesz index

Let  $(A, P, N)$  a Poisson-Nijenhuis Lie algebroid. For any  $x \in M$  consider the map  $N_x: A_x \rightarrow A_x$ . Recall that there exists a smallest integer  $k > 0$  such that the sequences

$$\text{Im } N_x \supseteq \text{Im } N_x^2 \supseteq \dots$$

and

$$\ker N_x \subseteq \ker N_x^2 \subseteq \dots$$

both stabilize at rank  $k$ . That is,

$$\text{Im } N_x^k = \text{Im } N_x^{k+1} = \dots, \quad \text{while } \text{Im } N_x^{k-1} \neq \text{Im } N_x^k,$$

and

$$\ker N_x^k = \ker N_x^{k+1} = \dots, \quad \text{while } \ker N_x^{k-1} \neq \ker N_x^k.$$

The integer  $k$  is called the **Riesz index** of  $N$  at  $x$ .

# The Reduced nondegenerate SN Lie algebroid

## Theorem 2

Let  $(A \rightarrow M, [\cdot, \cdot]_A, \rho_A, \Omega, N)$  be a symplectic-Nijenhuis Lie algebroid such that

- 1)  $N$  has constant Riesz index  $k$ .
- 2)  $\rho_A(B)$  and  $\mathcal{F}$  are regular foliations for  $B = \ker N^k$ .
- 3) For all  $x \in M$ ,  $a_x, a'_x \in L_{\mathcal{F}} \implies a_x - a'_x \in \ker(N_x^k)$ .

Then, we can induce a symplectic-Nijenhuis Lie algebroid structure  $([\cdot, \cdot]_{\tilde{A}}, \rho_{\tilde{A}}, \tilde{\Omega}, \tilde{N})$  on  $\tilde{A} = A/\mathcal{F} \longrightarrow \tilde{M} = M/\rho_A(\ker N^k)$  with  $\tilde{N}$  nondegenerate.

# The Reduced nondegenerate SN Lie algebroid

## Theorem 2

Let  $(A \rightarrow M, [\cdot, \cdot]_A, \rho_A, \Omega, N)$  be a symplectic-Nijenhuis Lie algebroid such that

- 1)  $N$  has constant Riesz index  $k$ .
- 2)  $\rho_A(B)$  and  $\mathcal{F}$  are regular foliations for  $B = \ker N^k$ .
- 3) For all  $x \in M$ ,  $a_x, a'_x \in L_{\mathcal{F}} \implies a_x - a'_x \in \ker(N_x^k)$ .

Then, we can induce a symplectic-Nijenhuis Lie algebroid structure  $([\cdot, \cdot]_{\tilde{A}}, \rho_{\tilde{A}}, \tilde{\Omega}, \tilde{N})$  on  $\tilde{A} = A/\mathcal{F} \longrightarrow \tilde{M} = M/\rho_A(\ker N^k)$  with  $\tilde{N}$  nondegenerate.

# The Reduced nondegenerate SN Lie algebroid

## Theorem 2

Let  $(A \rightarrow M, [\cdot, \cdot]_A, \rho_A, \Omega, N)$  be a symplectic-Nijenhuis Lie algebroid such that

- 1)  $N$  has constant Riesz index  $k$ .
- 2)  $\rho_A(B)$  and  $\mathcal{F}$  are regular foliations for  $B = \ker N^k$ .
- 3) For all  $x \in M$ ,  $a_x, a'_x \in L_{\mathcal{F}} \implies a_x - a'_x \in \ker(N_x^k)$ .

Then, we can induce a symplectic-Nijenhuis Lie algebroid structure  $([\cdot, \cdot]_{\tilde{A}}, \rho_{\tilde{A}}, \tilde{\Omega}, \tilde{N})$  on  $\tilde{A} = A/\mathcal{F} \longrightarrow \tilde{M} = M/\rho_A(\ker N^k)$  with  $\tilde{N}$  nondegenerate.

# The Reduced nondegenerate SN Lie algebroid

## Theorem 2

Let  $(A \rightarrow M, [\cdot, \cdot]_A, \rho_A, \Omega, N)$  be a symplectic-Nijenhuis Lie algebroid such that

- 1)  $N$  has constant Riesz index  $k$ .
- 2)  $\rho_A(B)$  and  $\mathcal{F}$  are regular foliations for  $B = \ker N^k$ .
- 3) For all  $x \in M$ ,  $a_x, a'_x \in L_{\mathcal{F}} \implies a_x - a'_x \in \ker(N_x^k)$ .

Then, we can induce a symplectic-Nijenhuis Lie algebroid structure  $([\cdot, \cdot]_{\tilde{A}}, \rho_{\tilde{A}}, \tilde{\Omega}, \tilde{N})$  on  $\tilde{A} = A/\mathcal{F} \longrightarrow \tilde{M} = M/\rho_A(\ker N^k)$  with  $\tilde{N}$  nondegenerate.

# The Reduced nondegenerate SN Lie algebroid

## Theorem 2

Let  $(A \rightarrow M, [\cdot, \cdot]_A, \rho_A, \Omega, N)$  be a symplectic-Nijenhuis Lie algebroid such that

- 1)  $N$  has constant Riesz index  $k$ .
- 2)  $\rho_A(B)$  and  $\mathcal{F}$  are regular foliations for  $B = \ker N^k$ .
- 3) For all  $x \in M$ ,  $a_x, a'_x \in L_{\mathcal{F}} \implies a_x - a'_x \in \ker(N_x^k)$ .

Then, we can induce a symplectic-Nijenhuis Lie algebroid structure  $([\cdot, \cdot]_{\widetilde{A}}, \rho_{\widetilde{A}}, \widetilde{\Omega}, \widetilde{N})$  on  $\widetilde{A} = A/\mathcal{F} \longrightarrow \widetilde{M} = M/\rho_A(\ker N^k)$  with  $\widetilde{N}$  nondegenerate.

## Reduction:Summary

$$(A \rightarrow M, [\cdot, \cdot]_A, \rho_A, P, N)$$

Poisson-Nijenhuis Lie algebroid

## Reduction:Summary

$$(A \rightarrow M, [\cdot, \cdot]_A, \rho_A, P, N)$$

Poisson-Nijenhuis Lie algebroid

$$\Downarrow \quad D = \rho_A(P^\sharp(A^*))$$

$$(A_L = P^\sharp(A^*)|_L \rightarrow L, [\cdot, \cdot]_{A_L}, \rho_{A_L}, \Omega_L, N_L)$$

symplectic-Nijenhuis Lie algebroid

## Reduction:Summary

$$(A \rightarrow M, [\cdot, \cdot]_A, \rho_A, P, N)$$

Poisson-Nijenhuis Lie algebroid

$$\Downarrow \quad D = \rho_A(P^\sharp(A^*))$$

$$(A_L = P^\sharp(A^*)|_L \rightarrow L, [\cdot, \cdot]_{A_L}, \rho_{A_L}, \Omega_L, N_L)$$

symplectic-Nijenhuis Lie algebroid

$$\Downarrow \quad \mathcal{F} = \{X^c + Y^\vee/X, Y \in \Gamma(\ker N_I^k)\}$$

$$(\widetilde{A} = A_L/\mathcal{F} \rightarrow \widetilde{L} = L/\rho_{A_L}(\ker N_L^k), [\cdot, \cdot]_{\widetilde{A}}, \rho_{\widetilde{A}}, \Omega_{\widetilde{A}}, N_{\widetilde{A}})$$

symplectic-Nijenhuis Lie algebroid with  $N_{\tilde{A}}$  nondegenerate

# References

- F. Magri, C. Morosi, *A geometrical characterization of integrable Hamiltonian systems through the theory of Poisson-Nijenhuis manifolds*, Quaderno S 19, University of Milan (1984).
- Y. Kosmann-Schwarzbach and F. Magri, *Poisson-Nijenhuis structures*, Ann. Inst. Henri Poincaré **53** (1990), 35–81.
- C. M. Marle, J. Nunes da Costa, *Reduction of bi-Hamiltonian manifolds and recursion operators*, DGA (Brno, 1995), 523–538, Masaryk Univ., Brno, 1996.
- J. Grabowski and P. Urbanski, *Lie algebroids and Poisson-Nijenhuis structures*, Rep. Math. Phys. **40** (1997) 195–208.
- R. Caseiro, *Modular Classes of PoissonNijenhuis Lie Algebroids*, Lett. Math. Phys. **80** (2007) 223–238.