

Products in Jacobi Geometry

LUCA VITAGLIANO



University of Salerno, Italy

(based on joint work in progress with: J. SCHNITZER)

International Conference on Poisson Geometry
IMPA, Rio de Janeiro, December 09–13, 2019

(M_1, ω_1) and (M_2, ω_2) symplectic manifolds

\Downarrow

$(M_1 \times M_2, \omega_1 + \omega_2)$ a symplectic manifold

Lagrangian calculus

(M_1, π_1) and (M_2, π_2) Poisson manifolds

\Downarrow

$(M_1 \times M_2, \pi_1 + \pi_2)$ a Poisson manifold

Coisotropic calculus

Weinstein Splitting

Every Poisson manifold (M, π) is locally of the form

$$(N \times O, \pi_N + \omega_O^{-1})$$

with (N, π_N) having a 0-dimensional leaf.

Remarks

- ▶ $(M_1 \times M_2, \pi_1 + \pi_2)$ is *not* a categorical product;
- ▶ products of Dirac manifolds are Dirac;
- ▶ coisotropic/isotropic calculus for Dirac manifolds [UCHINO 2004];
- ▶ Splitting for Dirac manifolds [BLOHMANN 2017].

Products of Contact Manifolds?

Contact Geometry:

odd dimensional analogue of Symplectic Geometry.

Contact structure:

maximally non-integrable hyperplane distribution $H \subset TM$.

\rightsquigarrow Contact manifolds are odd-dimensional.

(M_1, H_1) and (M_2, H_2) contact manifolds, but...

$M_1 \times M_2$ *cannot be a contact manifold.*

A solution within (Dirac-)Jacobi Geometry via *Homogenization!*

- Contact Geometry revisited
- A Symplectic-to-Contact dictionary
- Homogenization
- Products[!] of Dirac-Jacobi manifolds
- Products and splittings

Derivations

$E \rightarrow M$ a vector bundle.

Derivation of E :

$$\Delta : \Gamma(E) \rightarrow \Gamma(E)$$

- Δ is \mathbb{R} -linear,
- *Leibniz rule*: $\Delta(fe) = X(f)e + f\Delta(e)$.

$X =: \sigma(\Delta)$ is the *symbol*.

Remarks

- ▶ Derivations are sections of the *gauge algebroid*: $DE \rightrightarrows M$;
- ▶ DE acts tautologically on E ;
- ▶ the de Rham complex $(\Omega^\bullet(DE, E), d_{DE})$ is acyclic;
- ▶ $E = L$ a line bundle $\Rightarrow DL = \text{Hom}(\mathfrak{J}^1 L, L)$ and $\mathfrak{J}^1 L = \text{Hom}(DL, L)$.

Symplectic Atiyah Forms

(M, H) a contact manifold,
 $L = TM/H$ the normal bundle,
 $\theta : TM \rightarrow L$ the projection.

$$\left\{ \begin{array}{l} \text{Contact structures } H \\ \text{with } TM/H = L \end{array} \right\} \rightleftharpoons \left\{ \begin{array}{l} \text{closed, non-degenerate} \\ \omega \in \Omega^2(DL, L) \end{array} \right\}$$
$$H \mapsto d_{DL}(\theta \circ \sigma)$$
$$\sigma(\ker \iota_H \omega) \leftarrow \omega$$

Symplectic-to-Contact Dictionary:

- ▶ $C^\infty(M) \rightsquigarrow \Gamma(L)$
- ▶ $TM \rightsquigarrow DL$
- ▶ $T^*M \rightsquigarrow \mathfrak{J}^1 L$

Jacobi Manifolds

Symplectic-to-Contact Dictionary:

- ▶ *Poisson* \rightsquigarrow *Jacobi*;
- ▶ *Dirac* \rightsquigarrow *Dirac-Jacobi*;
- ▶ *Generalized complex* \rightsquigarrow *Generalized contact*;

Jacobi manifold: manifold M with a line bundle $L \rightarrow M$ and

$$\{-, -\} : \Gamma(L) \times \Gamma(L) \rightarrow \Gamma(L) \quad (\text{Jacobi bracket})$$

- a Lie bracket,
- a bi-derivation.

Jacobi Geometry

A Jacobi bracket can be seen as $J : \wedge^2 \mathfrak{J}^1 L \rightarrow L$ (*Jacobi tensor*).

When $L = \mathbb{R}_M$ then $\mathfrak{J}^1 L = T^*M \oplus \mathbb{R}_M$ and

$$J \equiv (\Lambda, E) \in \mathfrak{X}^2(M) \oplus \mathfrak{X}(M)$$

$$J^\# : \mathfrak{J}^1 L \rightarrow DL$$

$\text{Im}(\sigma \circ J^\#) \subset TM$ integrable distribution \rightsquigarrow characteristic foliation.

Remark

- ▶ Jacobi manifolds possess two kinds of leaves:
 - odd-dimensional \rightsquigarrow contact leaves;
 - even-dimensional \rightsquigarrow lcs leaves.

J invertible $\Leftrightarrow \omega = J^{-1}$ symplectic Atiyah form $\Leftrightarrow H$ contact structure.

Dirac-Jacobi Manifolds

$\mathbb{T}M = TM \oplus T^*M \rightsquigarrow \mathbb{D}L = DL \oplus \mathfrak{J}^1 L$: *omni-Lie algebroid* [CHEN, LIU 2010]

Structures on $\mathbb{D}L$:

$$\langle\langle (\Delta, \phi), (\nabla, \psi) \rangle\rangle = \langle \Delta, \psi \rangle + \langle \nabla, \phi \rangle,$$

$$\llbracket (\Delta, \phi), (\nabla, \psi) \rrbracket = ([\Delta, \nabla], \mathcal{L}_\Delta \psi - \iota_\nabla d_{DL} \phi).$$

Dirac-Jacobi manifold: [WADE 2000][V 2018]

manifold M with a line bundle L and a subbundle $\mathfrak{L} \subset \mathbb{D}L$

- \mathfrak{L} *maximally isotropic*,
- \mathfrak{L} *involutive*.

Remarks

- ▶ There are backward and forward Dirac-Jacobi maps,
- ▶ Dirac-Jacobi manifolds possess two kinds of leaves:
 - pre-contact leaves;
 - lc pre-s leaves.

Examples

- $DL \subset \mathbb{D}L$ and $\mathfrak{J}^1 L \subset \mathbb{D}L$,
- Jacobi brackets $\{-, -\} \equiv J$ ($\mathfrak{L}_J = \text{graph}(J^\#)$),
- pre-contact structures (= hyperplane distributions),
- locally conformal pre-symplectic structures,
- Dirac structures,
- *homogeneous Poisson structures*.

From Homogeneous Poisson to Dirac-Jacobi

Homogeneous Poisson manifold (Infinitesimal):

manifold P with (Z, π)

- Z a vector field such that,
- π a Poisson bivector

$$\mathcal{L}_Z \pi = -\pi$$

$$\left\{ \begin{array}{l} \text{Homogeneous Poisson} \\ \text{structures on } P \end{array} \right\} \rightarrow \left\{ \begin{array}{l} \text{Dirac-Jacobi} \\ \text{structures on } \mathbb{R}P \end{array} \right\} \quad [\text{WADE 2000}]$$
$$(Z, \pi) \mapsto \mathfrak{L}_{(Z, \pi)}$$

$$\mathfrak{L}_{(Z, \pi)} = \left\{ \left(h - hZ + \pi^\sharp(\eta), \eta \otimes 1 + \eta(Z) \cdot j^1 1 \right) : h \in \mathbb{R}, \eta \in T^*P \right\}$$

From Contact to Homogeneous Symplectic

$$\{\text{Contact manifolds}\} \Leftrightarrow \{\text{Homogeneous symplectic manifolds}\}$$

Homogeneous symplectic manifold (Finite):

manifold P with (h, ω)

- $h : \mathbb{R}^* \times P \rightarrow P$ a free and proper action,
- ω a symplectic form such that

$$h_r^*(\omega) = r \cdot \omega$$

$$\{\text{Line bundles}\} \Leftrightarrow \{\text{Principal } \mathbb{R}^*\text{-bundles}\}$$

$$L \mapsto \tilde{L} = L^* \setminus 0$$

$$(P \times \mathbb{R})/\mathbb{R}^* \leftarrow P$$

Symplectization

$L \rightarrow M$ a line bundle,
 $\tilde{L} \xrightarrow{p} M$ the principal \mathbb{R}^* -bundle.

$$\begin{array}{ccc} \mathbb{R}\tilde{L} & \longrightarrow & L \\ \downarrow \lrcorner & & \downarrow \\ \tilde{L} & \xrightarrow{p} & M \end{array} \quad \begin{array}{ccc} T\tilde{L} & \longrightarrow & DL \\ \downarrow \lrcorner & & \downarrow \\ \tilde{L} & \xrightarrow{p} & M \end{array} \quad \begin{array}{ccc} T^*\tilde{L} & \longrightarrow & \mathfrak{J}^1 L \\ \downarrow \lrcorner & & \downarrow \\ \tilde{L} & \xrightarrow{p} & M \end{array}$$

contact structure H with $TM/H = L$

$\omega \in \Omega^2(DL, L)$ symplectic Atiyah 2-form

$\tilde{\omega} = p^*\omega$ homogeneous symplectic structure on \tilde{L}

$$\begin{aligned} \{\text{Contact manifolds}\} &\Leftrightarrow \{\text{Homogeneous symplectic manifolds}\} \\ (M, H) &\mapsto (\tilde{L}, \tilde{\omega}) \end{aligned}$$

Poisson-ization and Dirac-ization

$L \rightarrow M$ a line bundle,
 $\tilde{L} \xrightarrow{p} M$ the principal \mathbb{R}^* -bundle.

$$\begin{aligned} \{\text{Jacobi manifolds}\} &\Leftrightarrow \{\text{Homogeneous Poisson manifolds}\} \\ (M, L, \{-, -\} \equiv J) &\mapsto (\tilde{L}, \tilde{J} := p^*J) \end{aligned}$$

$$\begin{aligned} \{\text{Dirac-Jacobi manifolds}\} &\Leftrightarrow \{\text{Homogeneous Dirac manifolds}\} \\ (M, L, \mathfrak{L} \subset \mathbb{D}L) &\mapsto (\tilde{L}, \tilde{\mathfrak{L}} := p^*\mathfrak{L} \subset \mathbb{T}\tilde{L}) \end{aligned}$$

Homogeneous Poisson structure (Finite): $h_r^*(\pi) = r^{-1} \cdot \pi$,

Homogeneous Dirac structure: $(Th_r \oplus r \cdot T^*h_{r^{-1}})(\mathfrak{L}) = \mathfrak{L}$.

Products[!] of Line Bundles

LB = {line bundles, fiber-wise invertible line bundle maps}.

Lemma

LB has products.

Proof. $L_1 \rightarrow M_1, L_2 \rightarrow M_2$ line bundles $\rightsquigarrow \tilde{L}_1, \tilde{L}_2$ principal \mathbb{R}^* -bundles.

$$(\tilde{L}_1 \times \tilde{L}_2, h \times h)$$

is a product in the category of principal \mathbb{R}^* -bundles.

$$\tilde{L}_1 \times \tilde{L}_2 \rightsquigarrow L_1 \times^! L_2 \rightarrow M_1 \times^! M_2$$

and $L_1 \times^! L_2$ is a product in **LB**.

Products[!] of Line Bundles

$$\blacktriangleright M_1 \times^! M_2 = \text{Iso}(L_1, L_2) = \left\{ \varphi : (L_1)_x \xrightarrow{\cong} (L_2)_y \mid (x, y) \in M_1 \times M_2 \right\},$$

$$\blacktriangleright L_1 \times^! L_2 = (M_1 \times^! M_2) \times_{M_1} L_1 \cong (M_1 \times^! M_2) \times_{M_2} L_2.$$

$$\begin{array}{ccccc}
 L_1 & \longleftarrow & L_1 \times^! L_2 & \longrightarrow & L_2 \\
 \downarrow & & \downarrow & & \downarrow \\
 M_1 & \longleftarrow & M_1 \times^! M_2 & \longrightarrow & M_2 \\
 \parallel & & \downarrow & & \parallel \\
 M_1 & \longleftarrow & M_1 \times M_2 & \longrightarrow & M_2
 \end{array}$$

$$\left\{ \begin{array}{c} \text{sections} \\ M_1 \times^! M_2 \\ \downarrow \uparrow \sigma \\ M_1 \times M_2 \end{array} \right\} \Leftrightarrow \left\{ \begin{array}{c} \text{diagrams} \\ L_1 \longleftarrow L \longrightarrow L_2 \\ \downarrow \quad \downarrow \quad \downarrow \\ M_1 \longleftarrow M_1 \times M_2 \longrightarrow M_2 \end{array} \right\}$$

$$\sigma \mapsto L = \sigma^* (L_1 \times^! L_2)$$

Products[!] of Dirac-Jacobi Manifolds

Lemma

(M_i, L_i) Dirac-Jacobi, $i = 1, 2 \implies (M_1 \times^! M_2, L_1 \times^! L_2)$ Dirac-Jacobi.

Proof.

$$\begin{aligned} & (M_1, L_1, \mathfrak{L}_1), (M_2, L_2, \mathfrak{L}_2) \text{ Dirac-Jacobi manifolds} \\ & \quad \Downarrow \\ & (\tilde{L}_1, h, \tilde{\mathfrak{L}}_1), (\tilde{L}_2, h, \tilde{\mathfrak{L}}_2) \text{ homogeneous Dirac manifolds} \\ & \quad \Downarrow \\ & (\tilde{L}_1 \times \tilde{L}_2, h \times h, \tilde{\mathfrak{L}}_1 \times \tilde{\mathfrak{L}}_2) \\ & \quad \Downarrow \\ & (M_1 \times^! M_2, L_1 \times^! L_2, \mathfrak{L}_1 \times^! \mathfrak{L}_2) \end{aligned}$$

The Product[!] Dirac-Jacobi Structure

$$\begin{array}{ccccc}
 \mathfrak{L}_1 & & \mathfrak{L}_1 \times^! \mathfrak{L}_2 = p_1^! \mathfrak{L}_1 \star p_2^! \mathfrak{L}_2 & & \mathfrak{L}_2 \\
 L_1 \longleftarrow & & L_1 \times^! L_2 \longrightarrow & & L_2 \\
 \downarrow & & \downarrow & & \downarrow \\
 M_1 \xleftarrow{p_1} & & M_1 \times^! M_2 \xrightarrow{p_2} & & M_2 \\
 \parallel & & \downarrow & & \parallel \\
 M_1 \longleftarrow & & M_1 \times M_2 \longrightarrow & & M_2
 \end{array}$$

$$p^! \mathfrak{L} = \{(\Delta, p^* \phi) : (p_* \Delta, \phi) \in \mathfrak{L}\},$$

$$\mathfrak{L} \star \mathfrak{L}' = \{(\Delta, \phi + \phi') : (\Delta, \phi) \in \mathfrak{L}, (\Delta, \phi') \in \mathfrak{L}'\}.$$

Remarks

- ▶ contact $\times^!$ contact = contact,
- ▶ pre-contact $\times^!$ pre-contact = pre-contact,
- ▶ Jacobi $\times^!$ Jacobi = Jacobi.

Splittings of Jacobi Manifolds

$(M, L, \{-, -\} \equiv J)$ a Jacobi manifold.

Theorem^[DAZORD, LICHNEROWICZ, MARLE 1991]

Let $x_0 \in M$ be a point in a **contact leaf** \mathcal{O} .

- 1 Every *small complementary transversal* $N \ni x_0$ possesses a homogeneous Poisson structure (Z_N, π_N) such that $Z_N|_{x_0} = \pi_N|_{x_0} = 0$;
- 2 There is a local Jacobi isomorphism

$$(M, L, J) \cong (N \times \mathbb{R}^{odd}, \mathbb{R}_{N \times \mathbb{R}^{odd}}, J^\times)$$

where $J^\times \equiv (\Lambda^\times, E^\times)$ and

$$\Lambda^\times = \Lambda_{can} + \pi_N + E_{can} \wedge Z_N \quad \text{and} \quad E^\times = E_{can}.$$

Splittings of Jacobi Manifolds

$(M, L, \{-, -\} \equiv J)$ a Jacobi manifold.

Theorem^[DAZORD, LICHNEROWICZ, MARLE 1991]

Let $x_0 \in M$ be a point in a **lcs leaf** \mathcal{O} .

- 1 Every *small complementary transversal* $N \ni x_0$ possesses a Jacobi structure $J_N \equiv (\Lambda_N, E_N)$ such that $J_N|_{x_0} = 0$;
- 2 There is a local Jacobi isomorphism

$$(M, L, J) \cong (N \times \mathbb{R}^{\text{even}}, \mathbb{R}_{N \times \mathbb{R}^{\text{even}}}, J^\times)$$

where $J^\times \equiv (\Lambda^\times, E^\times)$ and

$$\Lambda^\times = \Lambda_N + \pi_{\text{can}} + E_N \wedge Z_{\text{can}} \quad \text{and} \quad E^\times = E_N.$$

Splittings of Jacobi Manifolds

The DLM splittings can be written in terms of products:

contact leaf: \exists a section $\sigma : N \times \mathbb{R}^{odd} \rightarrow N \times^! \mathbb{R}^{odd}$

$$\begin{aligned}\mathbb{R}_{N \times \mathbb{R}^{odd}} &= \sigma^* \left(\mathbb{R}_N \times^! \mathbb{R}_{\mathbb{R}^{odd}} \right) \\ \mathfrak{L}_{(\Lambda^\times, E^\times)} &= \sigma^! \left(\mathfrak{L}_{(Z_N, \pi_N)} \times^! \mathfrak{L}_{(\Lambda_{can}, E_{can})} \right)\end{aligned}$$

lcs leaf: \exists a section $\sigma : N \times \mathbb{R}^{even} \rightarrow N \times^! \mathbb{R}^{even}$

$$\begin{aligned}\mathbb{R}_{N \times \mathbb{R}^{even}} &= \sigma^* \left(\mathbb{R}_N \times^! \mathbb{R}_{\mathbb{R}^{even}} \right) \\ \mathfrak{L}_{(\Lambda^\times, E^\times)} &= \sigma^! \left(\mathfrak{L}_{(\Lambda_N, E_N)} \times^! \mathfrak{L}_{(Z_{can}, \pi_{can})} \right)\end{aligned}$$

Splittings of Dirac-Jacobi Manifolds

(M, L, \mathfrak{L}) a Dirac-Jacobi manifold.

Theorem^[SCHNITZER 2019]

Let $x_0 \in M$ be a point in a pre-contact leaf \mathcal{O} .

- 1 Every *small complementary transversal* $N \ni x_0$ possesses a homogeneous Poisson structure (Z_N, π_N) such that $Z_N|_{x_0} = \pi_N|_{x_0} = 0$;
- 2 There is
 - a local section $\sigma : N \times \mathcal{O} \rightarrow N \times^! \mathcal{O}$ and
 - a local Dirac-Jacobi isomorphism up to a B -field

$$(M, L, \mathfrak{L}) \cong^B \left(N \times \mathcal{O}, \sigma^* \left(L|_N \times^! L|_{\mathcal{O}} \right), \sigma^! \left(\mathfrak{L}_{(Z_N, \pi_N)} \times^! DL|_{\mathcal{O}} \right) \right)$$

Splittings of Dirac-Jacobi Manifolds

(M, L, \mathfrak{L}) a Dirac-Jacobi manifold.

Theorem^[SCHNITZER 2019]

Let $x_0 \in M$ be a point in a **lc pre-s leaf** \mathcal{O} .

- 1 Every *small complementary transversal* $N \ni x_0$ possesses a Jacobi structure J_N such that $J_N|_{x_0} = 0$;
- 2 There is
 - a local section $\sigma : N \times \mathcal{O} \rightarrow N \times^! \mathcal{O}$ and
 - a local Dirac-Jacobi isomorphism up to a B -field

$$(M, L, \mathfrak{L}) \cong^B \left(N \times \mathcal{O}, \sigma^* \left(L|_N \times^! L|_{\mathcal{O}} \right), \sigma^! \left(\mathfrak{L}_{J_N} \times^! DL|_{\mathcal{O}} \right) \right)$$

- A. J. BRUCE, K. GRABOWSKA, AND J. GRABOWSKA,
Remarks on Contact and Jacobi Geometry,
SIGMA **13** (2017), 059, 22 pages.
- A. WADE,
Conformal Dirac Structures,
Lett. Math. Phys. **53** (2000), 331–348.
- L. V.,
Dirac-Jacobi Bundles,
J. Symplectic Geom. **16** (2018), 485–561.
- J. SCHNITZER,
*Normal Forms for Dirac-Jacobi Bundles
and Splitting Theorems for Jacobi Structures*,
arXiv:1901.00207.
- L. V., AND J. SCHNITZER,
Products in Jacobi Geometry,
in progress.

Thank you!